

Course program of Teoria delle Funzioni 1

Academic Year 2012-2013

P.D. Lamberti

Almost all material presented during this lecture course can be found in [1].

1. Preliminaries and notation¹. Basic facts from the theory of L^p -spaces in open subsets of \mathbb{R}^N . Minkowski's inequality for integrals. Convolution in \mathbb{R}^N . Young's inequality for convolution. Generalized Young's inequality for convolution (without proof). Mollifiers. Properties of mollifiers: pointwise convergence, uniform convergence, convergence in L^p . Density of $C_c^\infty(\Omega)$ in $L^p(\Omega)$ for $p \in [1, \infty[$. Fundamental Lemma of the Calculus of Variations.

2. Weak derivatives. Motivation: integration by parts in \mathbb{R}^N and weak formulation of a differential problem (the case of the Poisson problem for the Laplace operator). Definition of weak derivatives via integration by parts. Equivalent definitions: definition of weak derivatives via approximation by means of regular functions, definition of weak derivatives in \mathbb{R} via absolutely continuous functions. Weak differentiation under integral sign. Weak derivatives and convolution. Existence of intermediate weak derivatives.

3. Sobolev Spaces. Definition of the Sobolev Space $W^{l,p}(\Omega)$ and its variants $\widetilde{W}^{l,p}(\Omega)$, $w^{l,p}(\Omega)$. Completeness. Basic examples: an example of a function in the Sobolev space $W^{1,p}(\Omega)$ which is unbounded in any neighborhood of any point². Equivalent norms. The Nikodym's domain³. The notion of differential dimension of a function space and the differential dimension of the Sobolev space $w^{l,p}(\mathbb{R}^N)$. Lipschitz continuous functions⁴: classic derivatives and weak derivatives, extensions of Lipschitz continuous functions, the Rademacher's Theorem.

4. Approximation theorems. Density of $C_c^\infty(\mathbb{R}^N)$ in $W^{l,p}(\mathbb{R}^N)$ for $p \in [1, \infty[$. The space $W_0^{l,p}(\Omega)$. Partition of unity and density⁵ of $C^\infty(\Omega) \cap W^{l,p}(\Omega)$ in $W^{l,p}(\Omega)$ for $p \in [1, \infty[$. Counterexamples to the density of $C^\infty(\bar{\Omega})$ in $W^{l,p}(\Omega)$. An important consequence: characterization of weak derivatives via functions which are absolutely continuous in almost all lines parallel to the coordinate axes.

5. Integral representations. Star-shaped domains with respect to a ball and domains satisfying the cone condition. Taylor's formula in \mathbb{R}^N with remainder in integral form. Sobolev's integral representation formula. Consequences: pointwise estimates for functions and intermediate derivatives.

6. Embedding theorems. The notion of embedding. Continuous embeddings between Sobolev spaces are equivalent to the corresponding inclusions. Sobolev's embedding Theorem for the space $W^{l,p}(\Omega)$ into $C_b(\Omega)$ (the case $lp > N$).

¹For this chapter, we refer also to [4]

²For this example, we refer to [2]

³For this example, we refer to [5]

⁴For this part, we refer to [3]

⁵For this and the rest of Chapter 4, we have followed the approach presented in [5]

Sobolev's embedding Theorem for the space $W^{l,p}(\Omega)$ into $L^{q^*}(\Omega)$ where q^* is the Sobolev's exponent (the case $lp < N$): the argument via Young's inequality and the argument via Hardy-Littlewood-Sobolev inequality. The exponent q^* cannot be improved. Sobolev's embedding Theorem for the the space $W^{l,p}(\Omega)$ into $L^q(\Omega)$ for any $q \in [p, \infty[$ (the case $lp = N$). Gagliardo's inequality (the case $p = 1$ without proof). Example of an open set for which the Sobolev's embedding doesn't hold (outer cusps). The Poincaré's inequality. Applications: existence, uniqueness and stability of weak solutions to the Poisson problem for the Laplace operator.

7. Estimates for intermediate derivatives. Classes of open sets: open sets with resolved and quasi-resolved boundaries, open sets with continuous and quasi-continuous boundaries, open sets with Lipschitz boundaries, open sets with C^l -boundaries. Relation between open sets with Lipschitz boundaries and open sets satisfying the cone condition. Estimates for intermediate derivatives (without proof). Sobolev's embedding Theorem in the general case $W^{l,p}(\Omega) \subset W^{m,q}(\Omega)$.

8. Compact embeddings, extensions, traces, and applications. The notion of compact operator. The Kolmogorov criterion for compactness in $L^p(\mathbb{R}^N)$ (without proof). The Rellich-Kondrakov Theorem (proof in the case of the embedding of $W_0^{l,p}(\Omega)$ into $L^p(\Omega)$ when Ω has finite measure). The extension Theorem (without proof). Application of the Rellich-Kondrakov Theorem to the Helmholtz equation: existence of the first eigenvalue of the Laplace operator with Dirichlet boundary conditions. The Trace Theorem⁶: existence of the trace operator with values in $L^p(\partial\Omega)$. Definition of Besov-Nikolskii spaces $B_p^l(\mathbb{R}^N)$ and characterization of traces via Besov-Nikolskii spaces $B_p^l(\partial\Omega)$ (without proof). Application of the Trace Theorem to the Dirichlet problem for the Laplace operator: existence of weak solutions.

References

- [1] V.I. Burenkov, *Sobolev spaces on domains*, B.G. Teubner, Stuttgart, 1998.
- [2] L.C. Evans, *Partial Differential Equations*, Graduate Studies in Mathematics Vol. 19, American Mathematical Society, Providence, Rhode Island, 1998.
- [3] L.C. Evans and R.F. Gariepy, *Measure theory and fine properties of functions*, CRC Press, London, 1992.
- [4] G.B. Folland, *Real Analysis. Modern techniques and their applications*, John Wiley & Sons, Inc., New York, 1999.
- [5] V.G. Maz'ya and S.V. Poborchii, *Differentiable functions on bad domains*, World Scientific Publishing, 1997.

⁶For this, we refer to [2]