

MATLIS CATEGORY EQUIVALENCES FOR A RING EPIMORPHISM

SILVANA BAZZONI AND LEONID POSITSSELSKI

ABSTRACT. Under mild assumptions, we construct the two Matlis additive category equivalences for an associative ring epimorphism $u: R \rightarrow U$. Assuming that the ring epimorphism is homological of flat/projective dimension 1, we discuss the abelian categories of u -comodules and u -contramodules and construct the recollement of unbounded derived categories of R -modules, U -modules, and complexes of R -modules with u -co/contramodule cohomology. Further assumptions allow to describe the third category in the recollement as the unbounded derived category of the abelian categories of u -comodules and u -contramodules. For commutative rings, we also prove that any homological epimorphism of projective dimension 1 is flat. Injectivity of the map u is not required.

CONTENTS

Introduction	1
1. First Additive Category Equivalence	4
2. Second Additive Category Equivalence	7
3. Abelian Categories of u -Comodules and u -Contramodules	12
4. The Endomorphism Ring of the Two-Term Complex $(R \rightarrow U)$	14
5. When is the Class of Torsion Modules Hereditary?	15
6. Triangulated Matlis Equivalence	17
7. Two Fully Faithful Triangulated Functors	19
8. Kronecker Quiver Example	21
References	27

INTRODUCTION

The aim of this paper is to develop the basics of the theory of comodules and contramodules for an associative ring epimorphism in the maximal natural generality, and for the purpose of future reference. Let us start this introduction with explaining what the words in the paper's title mean.

A *ring epimorphism* $u: R \rightarrow U$ is a homomorphism of associative rings such that for every pair of parallel ring homomorphisms $f, g: U \rightrightarrows V$ the equation $fu = gu$ implies $f = g$. Equivalently, a ring homomorphism u is an epimorphism if and only if the two induced maps $u \otimes \text{id}$ and $\text{id} \otimes u: U \rightrightarrows U \otimes_R U$ are equal to each other, if and only if one or both of the maps $u \otimes \text{id}$ and $\text{id} \otimes u$ are isomorphisms, and

if and only if the multiplication map $U \otimes_R U \rightarrow U$ is an isomorphism. Further equivalent conditions for a ring map u to be an epimorphism are that the functor of restriction of scalars $u_*: U\text{-mod} \rightarrow R\text{-mod}$ is fully faithful, or that the functor $u_*: \text{mod-}U \rightarrow \text{mod-}R$ is fully faithful [24, Section XI.1]. In a ring epimorphism $R \rightarrow U$, the ring U is commutative whenever the ring R is.

The history of what is known as Matlis category equivalences goes back to the paper of Harrison [10], where two equivalences between certain full additive subcategories of the category of abelian groups were constructed. The first equivalence was provided by the functor of tensor product with the abelian group \mathbb{Q}/\mathbb{Z} , with the inverse functor $\text{Hom}_{\mathbb{Z}}(\mathbb{Q}/\mathbb{Z}, -)$. The second equivalence was given by the pair of functors $\text{Tor}_1^{\mathbb{Z}}(\mathbb{Q}/\mathbb{Z}, -)$ and $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Q}/\mathbb{Z}, -)$.

In Matlis' memoir [14, Section 3], the setting was generalized as follows. Let R be a commutative domain, Q be its field of quotients, and $K = Q/R$ be the quotient R -module. Then there are two equivalences between certain full additive subcategories of the category of R -modules. The first equivalence is provided by the functor of tensor product with the R -module K , and the inverse functor is $\text{Hom}_R(K, -)$. The second equivalence is given by the pair of functors $\text{Tor}_1^R(K, -)$ and $\text{Ext}_R^1(K, -)$, which are mutually inverse in restriction to the respective subcategories. Moreover, in the book [15] Matlis extended the first one of his two category equivalences to the setting with an arbitrary commutative ring R and its total ring of quotients Q .

Let us mention two further generalizations of the Matlis category equivalences in two different directions, which appeared in the two recent papers [20, 6]. In the paper [20, Section 5], the two Matlis additive category equivalences were constructed for a localization $S^{-1}R$ of a commutative ring R with respect to a multiplicative subset $S \subset R$. Injectivity of the map $R \rightarrow S^{-1}R$ was not assumed, but the assumption that the projective dimension of the R -module $S^{-1}R$ does not exceed 1 was made. In the paper [6, Section 4], the first Matlis category equivalence was constructed for certain injective epimorphisms of noncommutative rings $R \rightarrow Q$, where Q is the localization of R with respect to a one-sided Ore subset of regular elements.

In this paper, we construct the first Matlis additive category equivalence for any ring epimorphism $u: R \rightarrow U$ such that $\text{Tor}_1^R(U, U) = 0$, and the second Matlis category equivalence for any u such that $\text{Tor}_1^R(U, U) = 0 = \text{Tor}_2^R(U, U)$. Let us emphasize that *neither* injectivity of u , *nor* any condition on the projective or flat dimension of the R -module U is required for these results. Commutativity of the rings R and U is not assumed, either.

Furthermore, assuming that U has projective dimension at most 1 as a left R -module and flat dimension at most 1 as a right R -module, we construct what was called the *triangulated Matlis equivalence* in [20]. However, unlike in [20], we do not deduce the Matlis equivalences between additive categories of modules from the triangulated equivalence, but prove them separately. This allows to obtain the extra generality mentioned above.

The key role is played by the full subcategories of what we call *u -comodules* and *u -contramodules* in $R\text{-mod}$. The former is defined as the full subcategory of all

left R -modules annihilated by the derived functor $\mathrm{Tor}_{0,1}^R(U, -)$, while the latter is the Geigle–Lenzing right $\mathrm{Ext}_R^{0,1}$ -perpendicular subcategory to U in the category of left R -modules. Under the assumptions of the flat/projective dimension of U not exceeding 1, these are abelian categories with exact inclusion functors into $R\text{-mod}$. In the respective assumptions, we show that the u -comodules form a Grothendieck abelian category, while the abelian category of u -contra-modules is locally presentable with a projective generator. We also discuss adjoint functors to the identity inclusions of these full subcategories into the category of left R -modules.

The triangulated Matlis equivalence is an equivalence between the (bounded or unbounded) derived category of complexes R -modules with u -comodule cohomology modules and the similar derived category of complexes of R -modules with u -contra-module cohomology modules. The *recollement* of triangulated Matlis equivalence identifies both these triangulated categories with the Verdier quotient category of the derived category $\mathrm{D}^*(R\text{-mod})$ by the image of the fully faithful functor of restriction of scalars $u_*: \mathrm{D}^*(U\text{-mod}) \rightarrow \mathrm{D}^*(R\text{-mod})$ for a homological ring epimorphism u ,

$$(1) \quad \mathrm{D}_{u\text{-co}}(R\text{-mod}) \cong \mathrm{D}^*(R\text{-mod})/u_*\mathrm{D}^*(U\text{-mod}) \cong \mathrm{D}_{u\text{-ctra}}(R\text{-mod}).$$

Under certain additional assumptions (which hold whenever, but not only when, u is injective) the exact embedding functors of the full subcategories of u -comodules and u -contra-modules, $R\text{-mod}_{u\text{-co}} \rightarrow R\text{-mod}$ and $R\text{-mod}_{u\text{-ctra}} \rightarrow R\text{-mod}$, induce fully faithful functors between the derived categories, identifying the leftmost and the rightmost categories in (1) with the derived categories of the abelian categories $R\text{-mod}_{u\text{-co}}$ and $R\text{-mod}_{u\text{-ctra}}$. Hence one obtains an equivalence between the two derived categories,

$$(2) \quad \mathrm{D}^*(R\text{-mod}_{u\text{-co}}) \cong \mathrm{D}^*(R\text{-mod}_{u\text{-ctra}}).$$

We should mention that, in the same assumptions as ours, the equivalence of derived categories (2) was obtained in [5, Corollary 4.4] as a particular case of a general result about derived decomposition of abelian categories. The general approach in [5] is based on the technique of complete Ext-orthogonal pair in abelian categories, which was introduced by Krause and Št'ovíček in [12] (see also [4]). The same argument as in the present paper, going back to [18] and [20], is used in [5] in order to prove that the triangulated functors induced by the embeddings of abelian subcategories are fully faithful. One difference between our approaches is that in the present paper we also obtain the equivalences (1) holding under weaker assumptions.

One of the main results of this paper is based on some recent results of Hrbek and Angeleri Hügel–Hrbek [11, 1]. We show that whenever $u: R \rightarrow U$ is a homological ring epimorphism and U is an R -module of projective dimension 1, it follows that U is a flat R -module. Generalizing Matlis' classical result, we also show that, under a mild assumption on an epimorphism of commutative rings $u: R \rightarrow U$, the ring \mathfrak{R} of endomorphisms of the complex $R \rightarrow U$ in the derived category of R -modules is commutative. Under certain assumptions, it follows that the ring of endomorphisms of the R -module $U/R = \mathrm{coker} u$ is commutative, too.

In the last section, we compute the full subcategories of u -comodules and u -contramodules for certain ring epimorphisms originating from the finite-dimensional noncommutative algebra R associated with the Kronecker quiver. We are grateful to Jan Šťovíček for the suggestion to consider this example.

Acknowledgment. The first-named author is partially supported by grants BIRD163492 and DOR1690814 of Padova University. The second-named author is supported by research plan RVO: 67985840.

1. FIRST ADDITIVE CATEGORY EQUIVALENCE

Let $u: R \rightarrow U$ be an epimorphism of associative rings (i. e., a ring homomorphism such that the multiplication map $U \otimes_R U \rightarrow U$ is an isomorphism of R - R -bimodules). Then one has $U \otimes_R D \cong D \cong \text{Hom}_R(U, D)$ for all left U -modules D , and the functor of restriction of scalars $U\text{-mod} \rightarrow R\text{-mod}$ is fully faithful. The similar assertions hold for the right modules. We will say that a certain R -module “is a U -module” if it belongs to the image of the functor of restriction of scalars.

We will use the simple notation U/R for the cokernel of the map $u: R \rightarrow U$. So U/R is an R - R -bimodule.

A left R -module M is called a u -comodule (or a left u -comodule) if

$$U \otimes_R M = 0 = \text{Tor}_1^R(U, M).$$

Similarly, a right R -module N is said to be a u -comodule (or a right u -comodule) if $N \otimes_R U = 0 = \text{Tor}_1^R(N, U)$.

A left R -module C is called a u -contramodule (or a left u -contramodule) if

$$\text{Hom}_R(U, C) = 0 = \text{Ext}_R^1(U, C).$$

By [8, Proposition 1.1], the class of all left u -comodules is closed under direct sums, cokernels of morphisms, and extensions in $R\text{-mod}$. The class of all left u -contramodules is closed under products, kernels of morphisms, and extensions.

Example 1.1. The following example explains the “comodules and contramodules” terminology. Let $R = k[x]$ be the ring of polynomials in one variable over a field k , let $U = k[x, x^{-1}]$ be ring of Laurent polynomials, and let $u: R \rightarrow U$ be the natural inclusion. So one obtains the ring U from R by inverting the single element x .

Let \mathcal{C} be the coalgebra over k such that the dual topological algebra \mathcal{C}^* is identified with the ring of formal power series $k[[x]]$. Then the full subcategory of u -comodules in $R\text{-mod}$ is equivalent to the category of comodules over the coalgebra \mathcal{C} , while the full subcategory of u -contramodules in $R\text{-mod}$ is equivalent to the category of \mathcal{C} -contramodules [17, Sections 1.3 and 1.6].

We will use the notation $\text{pd}_R E$ for the projective dimension of a left R -module E and $\text{fd}_R E$ for the flat dimension of a right R -module E .

Borrowing the terminology going back to Harrison [10] and Matlis [14], we will say that a left R -module A is u -torsion-free if it is an R -submodule of a left U -module,

or equivalently, if the map $A \rightarrow U \otimes_R A$ induced by the ring homomorphism u is injective. Similarly, we will say that a left R -module B is *u-h-divisible* if it is a quotient module of a left U -module, or equivalently, if the map $\text{Hom}_R(U, B) \rightarrow B$ induced by u is surjective.

Clearly, the class of all u -torsion-free left R -modules is closed under subobjects, direct sums, and products in $R\text{-mod}$. Any left R -module A has a unique maximal u -torsion-free quotient module, which can be computed as the image of the natural R -module morphism $A \rightarrow U \otimes_R A$. The class of all u -h-divisible left R -modules is closed under quotients, direct sums, and products. Any left R -module B has a unique maximal u -h-divisible submodule, which can be computed as the image of the natural R -module morphism $\text{Hom}_R(U, B) \rightarrow B$.

A left R -module A is said to be *u-torsion* if its maximal u -torsion-free quotient module vanishes, or equivalently, if $U \otimes_R A = 0$. Indeed, the U -module $U \otimes_R A$ is always generated by the image of the map $u \otimes \text{id}_A: A \rightarrow U \otimes_R A$; hence if the image of $u \otimes \text{id}_A$ vanishes, then so does the whole module $U \otimes_R A$. A left R -module B is said to be *u-h-reduced* if its maximal u -h-divisible submodule vanishes, or equivalently, if $\text{Hom}_R(U, B) = 0$. Indeed, the map $\text{Hom}(u, \text{id}_B): \text{Hom}_R(U, B) \rightarrow B$ assigns to an R -module morphism $f: U \rightarrow B$ the element $f(1) \in B$. The action of U in the left R -module $\text{Hom}_R(U, B)$ is given by the rule $(vf)(w) = f(wv)$ for all $v, w \in U$. Hence if image of the map $\text{Hom}(u, \text{id}_B)$ vanishes, then $f(v) = (vf)(1) = \text{Hom}(u, \text{id}_B)(vf) = 0$ for all $f \in \text{Hom}_R(U, B)$ and $v \in U$, so $f = 0$.

Let us warn the reader that our terminology is slightly confusing: a left R -module with no nonzero u -torsion submodules does *not* need to be u -torsion-free (unless $\text{fd } U_R \leq 1$, as we will see below). Similarly, a left R -module with no u -h-reduced quotient modules does not need to be u -h-divisible (unless $\text{pd } U_R \leq 1$). The problem is that, unless such homological dimension conditions are imposed on the R - R -bimodule U or the ring homomorphism u , the classes of u -torsion-free and u -h-divisible left R -modules do not need to be closed under extensions.

The following theorem provides what appears to be the maximal natural generality for the first of the two classical *Matlis category equivalences* [14, Theorem 3.4], [15, Corollary 2.4] (going back to Harrison's [10, Proposition 2.1]).

Theorem 1.2. *Assume that $\text{Tor}_1^R(U, U) = 0$. Then the restrictions of the adjoint functors $M \mapsto \text{Hom}_R(U/R, M)$ and $C \mapsto (U/R) \otimes_R C$ are mutually inverse equivalences between the additive categories of u -h-divisible left u -comodules M and u -torsion-free left u -contramodules C .*

Before proceeding to prove the theorem, let us formulate and prove a lemma.

Lemma 1.3. *If $\text{Tor}_1^R(U, U) = 0$, then*

- (a) *for any left R -module M , the left R -module $\text{Hom}_R(U/R, M)$ is a u -torsion-free u -contramodule;*
- (b) *for any left R -module C , the left R -module $(U/R) \otimes_R C$ is a u -h-divisible u -comodule.*

Proof. Part (a): the left R -module $\text{Hom}_R(U/R, M)$ is u -torsion-free as an R -submodule of the left U -module $\text{Hom}_R(U, M)$. Furthermore, since $U \otimes_R U = U$, we have $(U/R) \otimes_R U = 0$, and therefore $\text{Hom}_R(U, \text{Hom}_R(U/R, M)) = 0$.

To show that $\text{Ext}_R^1(U, \text{Hom}_R(U/R, M)) = 0$, one observes that our assumptions $U \otimes_R U = U$ and $\text{Tor}_1^R(U, U) = 0$ imply $\text{Tor}_1^R(U/R, U) = 0$, because the map $(R/\ker(u)) \otimes_R U \rightarrow U$ is an isomorphism.

For any associative rings R and S , left R -module L , S - R -bimodule E , and left S -module M such that $\text{Tor}_1^R(E, L) = 0$, there is a natural injective map of abelian groups

$$\text{Ext}_R^1(L, \text{Hom}_S(E, M)) \longrightarrow \text{Ext}_S^1(E \otimes_R L, M).$$

In particular, in the situation at hand $\text{Ext}_R^1(U, \text{Hom}_R(U/R, M))$ is a subgroup of $\text{Ext}_R^1((U/R) \otimes_R U, M) = 0$.

The proof of part (b) is dual-analogous. The left R -module $(U/R) \otimes_R C$ is u -h-divisible as a quotient R -module of the left U -module $U \otimes_R C$. Since $U \otimes_R (U/R) = 0$, we have $U \otimes_R (U/R) \otimes_R C = 0$.

For any associative rings R and S , right R -module B , R - S -bimodule E , and left S -module C such that $\text{Tor}_1^R(B, E) = 0$, there is a natural surjective map of abelian groups

$$\text{Tor}_1^S(B \otimes_R E, C) \longrightarrow \text{Tor}_1^R(B, E \otimes_S C).$$

In particular, in the situation at hand $\text{Tor}_1^R(U, (U/R) \otimes_R C)$ is a quotient group of $\text{Tor}_1^R(U \otimes_R (U/R), C) = 0$. For a more high-tech derived category/spectral sequence presentation of the same argument, see Lemmas 2.4–2.5 below. \square

Proof of Theorem 1.2. By Lemma 1.3, the functor $M \mapsto \text{Hom}_R(U/R, M)$ take u -h-divisible left u -comodules to u -torsion-free left u -contramodules and back (in fact, they take arbitrary left R -modules to left R -modules from these two classes). It remains to show that the restrictions of these functors to these two full subcategories in $R\text{-mod}$ are mutually inverse equivalences between them.

Let M be a u -h-divisible left u -comodule. We will show that the adjunction morphism $(U/R) \otimes_R \text{Hom}_R(U/R, M) \rightarrow M$ is an isomorphism. Since M is u -h-divisible, we have a natural short exact sequence of left R -modules

$$(3) \quad 0 \longrightarrow \text{Hom}_R(U/R, M) \longrightarrow \text{Hom}_R(U, M) \longrightarrow M \longrightarrow 0.$$

Since the left R -module $\text{Hom}_R(U/R, M)$ is u -torsion-free, we also have a natural short exact sequence of left R -modules

$$(4) \quad 0 \longrightarrow \text{Hom}_R(U/R, M) \longrightarrow U \otimes_R \text{Hom}_R(U/R, M) \longrightarrow U/R \otimes_R \text{Hom}_R(U/R, M) \longrightarrow 0.$$

Since M is a u -comodule, applying the functor $U \otimes_R -$ to the short exact sequence (3) produces an isomorphism $U \otimes_R \text{Hom}_R(U/R, M) \cong U \otimes_R \text{Hom}_R(U, M) = \text{Hom}_R(U, M)$.

Now the commutative diagram

$$\begin{array}{ccccc}
\mathrm{Hom}_R(U/R, M) & \longrightarrow & U \otimes_R \mathrm{Hom}_R(U/R, M) & \longrightarrow & U/R \otimes_R \mathrm{Hom}_R(U/R, M) \\
\parallel & & \downarrow \cong & & \downarrow \\
& & U \otimes_R \mathrm{Hom}_R(U, M) & & \\
& & \downarrow \cong & & \\
\mathrm{Hom}_R(U/R, M) & \longrightarrow & \mathrm{Hom}_R(U, M) & \longrightarrow & M
\end{array}$$

shows that we have a morphism from the short exact sequence (4) to the short exact sequence (3) that is the identity on the leftmost terms, an isomorphism on the middle terms, and the adjunction morphism on the rightmost terms. Therefore, the adjunction morphism is an isomorphism.

Let C be a u -torsion-free left u -contramodule. Let us show that the adjunction morphism $C \rightarrow \mathrm{Hom}_R(U/R, (U/R) \otimes_R C)$ is an isomorphism. Since C is u -torsion-free, we have a natural short exact sequence of left R -modules

$$(5) \quad 0 \longrightarrow C \longrightarrow U \otimes_R C \longrightarrow (U/R) \otimes_R C \longrightarrow 0.$$

Since the left R -module $(U/R) \otimes_R C$ is u -h-divisible, we also have a natural short exact sequence of left R -modules

$$(6) \quad 0 \longrightarrow \mathrm{Hom}_R(U/R, (U/R) \otimes_R C) \longrightarrow \mathrm{Hom}_R(U, (U/R) \otimes_R C) \longrightarrow (U/R) \otimes_R C \longrightarrow 0.$$

Since C is a u -contramodule, applying the functor $\mathrm{Hom}_R(U, -)$ to the short exact sequence (5) produces an isomorphism $U \otimes_R C = \mathrm{Hom}_R(U, U \otimes_R C) \cong \mathrm{Hom}_R(U, (U/R) \otimes_R C)$. Now the commutative diagram

$$\begin{array}{ccccc}
C & \longrightarrow & U \otimes_R C & \longrightarrow & (U/R) \otimes_R C \\
\downarrow & & \downarrow \cong & & \parallel \\
& & \mathrm{Hom}_R(U, U \otimes_R C) & & \\
& & \downarrow \cong & & \\
\mathrm{Hom}_R(U/R, (U/R) \otimes_R C) & \longrightarrow & \mathrm{Hom}_R(U, (U/R) \otimes_R C) & \longrightarrow & (U/R) \otimes_R C
\end{array}$$

shows that we have a morphism from the short exact sequence (5) to the short exact sequence (6) that is the identity on the rightmost terms, an isomorphism on the middle terms, and an adjunction morphism on the leftmost terms. Therefore, the adjunction morphism is an isomorphism. \square

2. SECOND ADDITIVE CATEGORY EQUIVALENCE

Let K^\bullet denote the two-term complex $R \rightarrow U$, with the term R placed in the cohomological degree -1 and the term U in the cohomological degree 0 . We will view

K^\bullet as an object of the bounded derived category of R - R -bimodules $\mathbf{D}^b(R\text{-mod-}R)$. So, there is a distinguished triangle

$$(7) \quad R \longrightarrow U \longrightarrow K^\bullet \longrightarrow R[1]$$

in the triangulated category $\mathbf{D}^b(R\text{-mod-}R)$.

Alternatively, the complex K^\bullet can be considered as an object of the bounded derived category of left R -modules $\mathbf{D}^b(R\text{-mod})$ endowed with a right action of the ring R by its derived category object endomorphisms, or as an object of the bounded derived category of right R -modules $\mathbf{D}^b(\text{mod-}R)$ endowed with a left action of R . Then (7) is viewed as a distinguished triangle in $\mathbf{D}^b(R\text{-mod})$ or $\mathbf{D}^b(\text{mod-}R)$.

By an abuse of notation, given a left R -module B , we will denote simply by

$$\text{Ext}_R^n(K^\bullet, B) = H^n(\mathbb{R}\text{Hom}_R(K^\bullet, B)) = \text{Hom}_{\mathbf{D}^b(R\text{-mod})}(K^\bullet, B[n])$$

the abelian group of all morphisms $K^\bullet \rightarrow B[n]$ in the derived category of left R -modules. The right action of R in the object $K^\bullet \in \mathbf{D}^b(R\text{-mod})$ induces a left R -module structure on the groups $\text{Ext}_R^n(K^\bullet, B)$.

Similarly, we set

$$\text{Tor}_n^R(K^\bullet, A) = H^{-n}(K^\bullet \otimes_R^{\mathbb{L}} A)$$

for any left R -module B . Here K^\bullet is viewed as an object of the bounded derived category of right R -modules for the purpose of computing the derived tensor product $K^\bullet \otimes_R^{\mathbb{L}} A$, and then the left action of R in the object $K^\bullet \in \mathbf{D}^b(\text{mod-}R)$ induces a left R -module structure on the groups $\text{Tor}_n^R(K^\bullet, A)$.

Lemma 2.1. *For every left R -module C , there are natural isomorphisms of left R -modules*

- (a) $\text{Tor}_n^R(K^\bullet, C) = 0 = \text{Ext}_R^n(K^\bullet, C)$ for $n < 0$;
- (b) $\text{Tor}_n^R(K^\bullet, C) = \text{Tor}_n^R(U, C)$ and $\text{Ext}_R^n(K^\bullet, C) = \text{Ext}_R^n(U, C)$ for all $n > 1$;
- (c) $\text{Tor}_0^R(K^\bullet, C) = (U/R) \otimes_R C$ and $\text{Ext}_R^0(K^\bullet, C) = \text{Hom}_R(U/R, C)$.

Proof. All the assertions follow immediately from the (co)homology long exact sequences obtained by applying the functors $\mathbb{R}\text{Hom}_R(-, C)$ and $-\otimes_R^{\mathbb{L}} C$ to the distinguished triangle (7). \square

Furthermore, for any left R -modules A and B there are five-term exact sequences of low-dimensional Tor and Ext induced by the distinguished triangle (7):

$$(8) \quad 0 \longrightarrow \text{Tor}_1^R(U, A) \longrightarrow \text{Tor}_1^R(K^\bullet, A) \\ \longrightarrow A \longrightarrow U \otimes_R A \longrightarrow \text{Tor}_0^R(K^\bullet, A) \longrightarrow 0$$

and

$$(9) \quad 0 \longrightarrow \text{Ext}_R^0(K^\bullet, B) \longrightarrow \text{Hom}_R(U, B) \longrightarrow B \\ \longrightarrow \text{Ext}_R^1(K^\bullet, B) \longrightarrow \text{Ext}_R^1(U, B) \longrightarrow 0.$$

Both (8) and (9) are exact sequences of left R -modules.

Borrowing the terminology of Matlis [14], we will say that a left R -module A is *u-special* if the map $A \rightarrow U \otimes_R A$ is surjective. Equivalently (in view of the exact

sequence (8) or Lemma 2.1(c)), this means that $\mathrm{Tor}_0^R(K^\bullet, A) = 0$. Similarly, a left R -module B is u -cospecial if the map $\mathrm{Hom}_R(U, B) \rightarrow B$ is injective. Equivalently (by the exact sequence (9) or Lemma 2.1(c)), this means that $\mathrm{Ext}_R^0(K^\bullet, B) = 0$.

The next lemma provides another characterization of u -special and u -cospecial modules.

Lemma 2.2. (a) *A left R -module A is u -special if and only if its maximal u -torsion-free quotient module is a U -module.*

(b) *A left R -module B is u -cospecial if and only if its maximal u -h-divisible submodule is a U -module.*

Proof. Part (b): if B is u -cospecial, then the morphism $\mathrm{Hom}_R(U, B) \rightarrow B$ is injective, so $\mathrm{Hom}_R(U, B)$ is the maximal u -h-divisible submodule of B . Conversely, if the maximal u -h-divisible submodule of B is a U -module D , then $\mathrm{Hom}_R(U/R, B) = \mathrm{Hom}_R(U/R, D) = 0$.

Part (a): if A is u -special, then the morphism $A \rightarrow U \otimes_R A$ is surjective, so $U \otimes_R A$ is the maximal u -torsion-free quotient module of A . Conversely, assume that the maximal u -torsion-free quotient module of A is a U -module D . Note that $\mathrm{Hom}_{\mathbb{Z}}(U/R, \mathbb{Q}/\mathbb{Z})$ is a u -torsion-free left R -module, because it is a submodule of the left U -module $\mathrm{Hom}_{\mathbb{Z}}(U, \mathbb{Q}/\mathbb{Z})$. Hence

$$\begin{aligned} \mathrm{Hom}_{\mathbb{Z}}((U/R) \otimes_R A, \mathbb{Q}/\mathbb{Z}) &\cong \mathrm{Hom}_R(A, \mathrm{Hom}_{\mathbb{Z}}(U/R, \mathbb{Q}/\mathbb{Z})) \\ &= \mathrm{Hom}_R(D, \mathrm{Hom}_{\mathbb{Z}}(U/R, \mathbb{Q}/\mathbb{Z})) \cong \mathrm{Hom}_{\mathbb{Z}}((U/R) \otimes_R D, \mathbb{Q}/\mathbb{Z}), \end{aligned}$$

and the last term is zero since D is a U -module. It follows that $(U/R) \otimes_R A = 0$. \square

The following theorem is our version of the second Matlis category equivalence [14, Theorem 3.8] (going back to Harrison's [10, Proposition 2.3]).

Theorem 2.3. *Assume that $\mathrm{Tor}_1^R(U, U) = 0 = \mathrm{Tor}_2^R(U, U)$. Then the restrictions of the functors $M \mapsto \mathrm{Ext}_R^1(K^\bullet, M)$ and $C \mapsto \mathrm{Tor}_1^R(K^\bullet, C)$ are mutually inverse equivalences between the additive categories of u -cospecial left u -comodules M and u -special left u -contramodules C .*

Before proving the theorem, we formulate two lemmas, which extend the result of Lemma 1.3.

Lemma 2.4. (a) *If $\mathrm{Tor}_1^R(U, U) = 0$, then the left R -module $\mathrm{Tor}_0^R(K^\bullet, A)$ is a u -comodule for any left R -module A .*

(b) *If $\mathrm{Tor}_1^R(U, U) = 0 = \mathrm{Tor}_R^2(U, U) = 0$, then the left R -module $\mathrm{Tor}_1^R(K^\bullet, A)$ is a u -comodule for any left R -module A such that $\mathrm{Tor}_0^R(K^\bullet, A) = 0$.*

(c) *If $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{fd} U_R \leq 1$, then the left R -module $\mathrm{Tor}_1^R(K^\bullet, A)$ is a u -comodule for any left R -module A .*

Proof. There is a spectral sequence

$$E_{pq}^2 = \mathrm{Tor}_p^R(U, \mathrm{Tor}_q^R(K^\bullet, A)) \implies E_{pq}^\infty = \mathrm{gr}_p \mathrm{Tor}_{p+q}^R(U \otimes_R^{\mathbb{L}} K^\bullet, A),$$

where $\mathrm{Tor}_n^R(U \otimes_R^{\mathbb{L}} K^\bullet, A) = H^{-n}(U \otimes_R^{\mathbb{L}} K^\bullet \otimes_R^{\mathbb{L}} A) = 0$ whenever $H^{-i}(U \otimes_R^{\mathbb{L}} K^\bullet) = 0$ for all $0 \leq i \leq n$. Since $U \otimes_R U = U$, the latter condition holds whenever $\mathrm{Tor}_i^R(U, U) = 0$ for all $1 \leq i \leq n$. Thus $E_{pq}^\infty = 0$ whenever $p + q \leq 1$ in the assumptions of part (a), whenever $p + q \leq 2$ in the assumptions of part (b), and for all $p, q \in \mathbb{Z}$ in the assumptions of part (c).

The differentials are $\partial_{pq}^r: E_{pq}^r \longrightarrow E_{p-r, q+r-1}^r$, $r \geq 2$. Now all the differentials involving $E_{0,0}^r$ and $E_{0,1}^r$ vanish for the dimension reasons, so $E_{0,0}^\infty = 0 = E_{0,1}^\infty$ implies $E_{0,0}^2 = 0 = E_{0,1}^2$. This proves part (a). Furthermore, the only possibly nontrivial differentials involving $E_{0,1}^r$ and $E_{1,1}^r$ are

$$\partial_{2,0}^2: E_{2,0}^2 \longrightarrow E_{0,1}^2 \quad \text{and} \quad \partial_{3,0}^2: E_{3,0}^2 \longrightarrow E_{1,1}^2.$$

When $\mathrm{Tor}_R^0(K^\bullet, A) = 0$, one has $E_{p,0}^2 = 0$ for all $p \in \mathbb{Z}$. When $\mathrm{fd} U_R \leq 1$, one has $E_{pq}^2 = 0$ for $p \geq 2$ and all q . In both cases, $E_{0,1}^\infty = 0 = E_{1,1}^\infty$ implies $E_{0,1}^2 = 0 = E_{1,1}^2$, proving parts (b) and (c). \square

Lemma 2.5. (a) *If $\mathrm{Tor}_1^R(U, U) = 0$, then the left R -module $\mathrm{Ext}_R^0(K^\bullet, B)$ is a u -contramodule for any left R -module B .*

(b) *If $\mathrm{Tor}_1^R(U, U) = 0 = \mathrm{Tor}_R^2(U, U) = 0$, then the left R -module $\mathrm{Ext}_R^1(K^\bullet, B)$ is a u -contramodule for any left R -module B such that $\mathrm{Tor}_R^0(K^\bullet, B) = 0$.*

(c) *If $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{pd}_R U \leq 1$, then the left R -module $\mathrm{Ext}_R^1(K^\bullet, B)$ is a u -contramodule for any left R -module B .*

Proof. Dual-analogous to Lemma 2.4 (and similar to [20, Lemma 1.7]). \square

Proof of Theorem 2.3. Let M be a u -cospecial left u -comodule. By Lemma 2.5(b), the left R -module $\mathrm{Ext}_R^1(K^\bullet, M)$ is a u -contramodule. Furthermore, the exact sequence (9) for the R -module M reduces to a four-term sequence

$$0 \longrightarrow \mathrm{Hom}_R(U, M) \longrightarrow M \longrightarrow \mathrm{Ext}_R^1(K^\bullet, M) \longrightarrow \mathrm{Ext}_R^1(U, M) \longrightarrow 0.$$

Denoting by E the image of the map $M \longrightarrow \mathrm{Ext}_R^1(K^\bullet, M)$, we have two short exact sequences of left R -modules $0 \longrightarrow \mathrm{Hom}_R(U, M) \longrightarrow M \longrightarrow E \longrightarrow 0$ and $0 \longrightarrow E \longrightarrow \mathrm{Ext}_R^1(K^\bullet, M) \longrightarrow \mathrm{Ext}_R^1(U, M) \longrightarrow 0$.

The assumptions that $U \otimes_R U = U$ and $\mathrm{Tor}_i^R(U, U) = 0$ for $i = 1$ and 2 imply that $U \otimes_R D = D$ and $\mathrm{Tor}_i^R(U, D) = 0$ for all left U -modules D and $i = 1, 2$. Hence (by Lemma 2.1 and the exact sequence (8) for the R -module D) we have $\mathrm{Tor}_i^R(K^\bullet, D) = 0$ for $-1 \leq i \leq 2$. In particular, this applies to the left U -modules $D = \mathrm{Hom}_R(U, M)$ and $\mathrm{Ext}_R^1(U, M)$.

Now from the long exact sequences of $\mathrm{Tor}_*^R(K^\bullet, -)$ related to our two short exact sequences of left R -modules we see that both the maps $\mathrm{Tor}_i^R(K^\bullet, M) \longrightarrow \mathrm{Tor}_i^R(K^\bullet, E) \longrightarrow \mathrm{Tor}_i^R(K^\bullet, \mathrm{Ext}_R^1(K^\bullet, M))$ are isomorphisms for $i = 0$ and 1 . In particular, $\mathrm{Tor}_0^R(K^\bullet, \mathrm{Ext}_R^1(K^\bullet, M)) \cong \mathrm{Tor}_0^R(K^\bullet, M) = (U/R) \otimes_R M = 0$, since $U \otimes_R M = 0$. Hence the left R -module $\mathrm{Ext}_R^1(K^\bullet, M)$ is u -special.

Furthermore, the map $\mathrm{Tor}_1^R(K^\bullet, M) \longrightarrow M$ in the short exact sequence (8) is an isomorphism, since $U \otimes_R M = 0 = \mathrm{Tor}_R^1(U, M)$. Thus we obtain a natural isomorphism $\mathrm{Tor}_1^R(K^\bullet, \mathrm{Ext}_R^1(K^\bullet, M)) \cong M$.

The dual-analogous argument shows that the left R -module $\mathrm{Tor}_1^R(K^\bullet, C)$ is a u -cospecial u -comodule for any u -special u -contramodule C , and provides a natural isomorphism $\mathrm{Ext}_R^1(K^\bullet, \mathrm{Tor}_1^R(K^\bullet, C)) \cong C$. One has to observe that $\mathrm{Hom}_R(U, D) = D$ and $\mathrm{Ext}_R^i(U, D) = 0$ for all left U -modules D and $i = 1, 2$, hence $\mathrm{Ext}_R^i(K^\bullet, D) = 0$ for $-1 \leq i \leq 2$, etc. \square

In the rest of this section we discuss how our theory simplifies and improves in the assumptions that the projective dimension of the left R -module U and/or the flat dimension of the right R -module U do not exceed 1.

Lemma 2.6. (a) *Assume that $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{fd} U_R \leq 1$. Then a left R -module A is u -torsion-free if and only if $\mathrm{Tor}_1^R(K^\bullet, A) = 0$.*

(b) *Assume that $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{pd}_R U \leq 1$. Then a left R -module B is u - h -divisible if and only if $\mathrm{Ext}_R^1(K^\bullet, B) = 0$.*

Proof. This is similar to [20, Lemma 5.1(b)]. Let us prove part (a). The “if” claim follows immediately from the exact sequence (8). To prove the “only if”, assume that A is u -torsion-free. Then the exact sequence (8) implies that the left R -module morphism $\mathrm{Tor}_1^R(U, A) \rightarrow \mathrm{Tor}_1^R(K^\bullet, A)$ is an isomorphism. Since $\mathrm{Tor}_1^R(K^\bullet, A)$ is a left u -comodule by Lemma 2.4(c) and $\mathrm{Tor}_1^R(U, A)$ is a left U -module, they can only be isomorphic when both of them vanish. \square

It is clear from the definition and Lemma 2.6(a) that, when $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{fd} U_R \leq 1$, the full subcategory of u -torsion-free R -modules is closed under extensions, subobjects, direct sums, and products. So u -torsion-free R -modules form the torsion-free class of a certain torsion pair in $R\text{-mod}$. The related torsion class is the class of all u -torsion R -modules, that is, all left R -modules A such that $U \otimes_R A = 0$.

Similarly, it is clear from the definition and Lemma 2.6(b) that, whenever $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{pd}_R U \leq 1$, the full subcategory of u - h -divisible R -modules is closed under extensions, quotients, direct sums and products. So u - h -divisible R -modules form the torsion class of a certain torsion theory in $R\text{-mod}$. The related torsion class is the class of all u - h -reduced R -modules, that is, all left R -modules B such that $\mathrm{Hom}_R(U, B) = 0$.

It is clear from the definition that the full subcategory of u -special left R -modules is closed under extensions, quotients, and direct sums. Hence it is the torsion class of a torsion pair in $R\text{-mod}$. When $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{fd} U_R \leq 1$, the related torsion-free class can be described as the class of all u -torsion-free u - h -reduced left R -modules.

Similarly, the full subcategory of u -cospecial left R -modules is closed under extensions, subobjects, direct sums, and products. Hence it is the torsion-free class of a torsion pair in $R\text{-mod}$. When $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{pd}_R U \leq 1$, the related torsion class can be described as the class of all u - h -divisible u -torsion left R -modules.

3. ABELIAN CATEGORIES OF u -COMODULES AND u -CONTRAMODULES

In this section, as in the previous one, $u: R \rightarrow U$ is an associative ring epimorphism. For most of the results, we will have to assume that u is a *homological* ring epimorphism, that is, $\mathrm{Tor}_i^R(U, U) = 0$ for $i \geq 1$. In fact, we will mostly have to assume either that the flat dimension of the right R -module U does not exceed 1 (when discussing left u -comodules), or that the projective dimension of the left R -module U does not exceed 1 (when considering left u -contramodules).

Let us denote the full subcategory of left u -comodules by $R\text{-mod}_{u\text{-co}} \subset R\text{-mod}$, and the full subcategory of left u -contramodules by $R\text{-mod}_{u\text{-ctra}} \subset R\text{-mod}$. For any left R -module C , we set $\Gamma_u(C) = \mathrm{Tor}_1^R(K^\bullet, C)$ and $\Delta_u(C) = \mathrm{Ext}_R^1(K^\bullet, C)$. The natural left R -module morphisms (occurring in the exact sequences (8–9)) are denoted by $\gamma_{u,C}: \Gamma_u(C) \rightarrow C$ and $\delta_{u,C}: C \rightarrow \Delta_u(C)$.

Proposition 3.1. *Assume that $\mathrm{fd} U_R \leq 1$. Then*

(a) *the full subcategory $R\text{-mod}_{u\text{-co}}$ is closed under the kernels, cokernels, extensions, and direct sums in $R\text{-mod}$. So $R\text{-mod}_{u\text{-co}}$ is an abelian category and the embedding functor $R\text{-mod}_{u\text{-co}} \rightarrow R\text{-mod}$ is exact;*

(b) *assuming also that $\mathrm{Tor}_1^R(U, U) = 0$, the functor $\Gamma_u: R\text{-mod} \rightarrow R\text{-mod}_{u\text{-co}}$ is right adjoint to the fully faithful embedding functor $R\text{-mod}_{u\text{-co}} \rightarrow R\text{-mod}$.*

Proof. Part (a) is a particular case of [8, Proposition 1.1] or [19, Theorem 1.2(b)]. To prove part (b), notice that $\Gamma_u(A) \in R\text{-mod}_{u\text{-co}}$ for any $A \in R\text{-mod}$ by Lemma 2.4(c). We have to show that for every left R -module A , every left u -comodule M , and an R -module morphism $M \rightarrow A$ there exists a unique R -module morphism $M \rightarrow \Gamma_u(A)$ making the triangle diagram $M \rightarrow \Gamma_u(A) \rightarrow A$ commutative.

Indeed, looking on the exact sequence (8), the composition $M \rightarrow A \rightarrow U \otimes_R A$ vanishes, since $U \otimes_R M = 0$. Now the obstruction to lifting the morphism $M \rightarrow A$ to a morphism $M \rightarrow \mathrm{Tor}_1^R(K^\bullet, A)$ lies in the group $\mathrm{Ext}_R^1(M, \mathrm{Tor}_1^R(U, A))$, and the obstruction to uniqueness of such a lifting lies in the group $\mathrm{Hom}_R(M, \mathrm{Tor}_1^R(U, A))$.

Generally, for any ring homomorphism $R \rightarrow U$, left R -module M , left U -module D , and an integer $n \geq 0$ such that $\mathrm{Tor}_i^R(U, M) = 0$ for $0 \leq i \leq n$, one has $\mathrm{Tor}_i^R(M, D) \cong \mathrm{Tor}_i^U(U \otimes_R M, D)$ and $\mathrm{Ext}_R^i(M, D) \cong \mathrm{Ext}_U^i(U \otimes_R M, D)$. In the situation at hand, $D = \mathrm{Tor}_1^R(U, A)$ is a left U -module, so any R -module morphism $M \rightarrow D$ vanishes, since $U \otimes_R M = 0$. Finally, we have $\mathrm{Ext}_R^1(M, D) = \mathrm{Ext}_U^1(U \otimes_R M, D) = 0$, since $\mathrm{Tor}_1^R(U, M) = 0$ and $U \otimes_R M = 0$. \square

Proposition 3.2. *Assume that $\mathrm{pd}_R U \leq 1$. Then*

(a) *the full subcategory $R\text{-mod}_{u\text{-ctra}}$ is closed under the kernels, cokernels, extensions, and products in $R\text{-mod}$. So $R\text{-mod}_{u\text{-ctra}}$ is an abelian category and the embedding functor $R\text{-mod}_{u\text{-ctra}} \rightarrow R\text{-mod}$ is exact;*

(b) *assuming also that $\mathrm{Tor}_1^R(U, U) = 0$, the functor $\Delta_u: R\text{-mod} \rightarrow R\text{-mod}_{u\text{-ctra}}$ is left adjoint to the fully faithful embedding functor $R\text{-mod}_{u\text{-ctra}} \rightarrow R\text{-mod}$.*

Proof. Part (a) is a particular case of [8, Proposition 1.1] or [19, Theorem 1.2(a)]. The proof of part (b) is based on Lemma 2.5(c) and dual-analogous to the proof of Proposition 3.1(b); cf. [20, Theorem 3.4]. \square

Lemma 3.3. *Assume that $\text{fd} U_R \leq 1$, $\text{pd}_R U \leq 1$, and $\text{Tor}_1^R(U, U) = 0$. Then*

(a) *for any u -h-divisible left R -module B , the left R -module $\Gamma_u(B)$ is also u -h-divisible;*

(b) *for any u -torsion-free left R -module A , the left R -module $\Delta_u(A)$ is also u -torsion-free.*

Proof. Let us prove part (a). Following Lemma 2.6(b), we have to check that $\text{Ext}_R^1(K^\bullet, \text{Tor}_1^R(K^\bullet, B)) = 0$. Since B is u -h-divisible, we have $\text{Tor}_0^R(K^\bullet, B) = (U/R) \otimes_R B = 0$, so the five-term exact sequence (8) reduces to a four-term sequence. Furthermore, $\text{Ext}_R^*(K^\bullet, D) = 0$ for any left U -module D . Thus it follows from (8) that $\text{Ext}_R^1(K^\bullet, \text{Tor}_1^R(K^\bullet, B)) = \text{Ext}_R^1(K^\bullet, B) = 0$ (cf. the proof of Theorem 2.3). The proof of part (b) is dual-analogous. \square

Lemma 3.4. *Assume that $\text{Tor}_1^R(U, U) = 0$ and $\text{fd} U_R \leq 1$. Then $R\text{-mod}_{u\text{-co}}$ is a Grothendieck abelian category. If J is an injective cogenerator of the abelian category $R\text{-mod}$, then $\Gamma_u(J)$ is an injective cogenerator of $R\text{-mod}_{u\text{-co}}$.*

Proof. By Proposition 3.1(a), the full subcategory $R\text{-mod}_{u\text{-co}}$ is closed under direct limits in $R\text{-mod}$; it is also an abelian category with an exact embedding functor $R\text{-mod}_{u\text{-co}} \rightarrow R\text{-mod}$. Hence the direct limit functors in $R\text{-mod}_{u\text{-co}}$ are exact, and it remains to show that this category has a set of generators.

By Proposition 3.1(b), the functor $\Gamma_u = \text{Tor}_1^R(K^\bullet, -)$ is right adjoint to the embedding functor $R\text{-mod}_{u\text{-co}} \rightarrow R\text{-mod}$. Viewed as a functor $R\text{-mod} \rightarrow R\text{-mod}$, the functor $\text{Tor}_1^R(K^\bullet, -)$ clearly preserves direct limits; hence it follows that the functor $\Gamma_u: R\text{-mod} \rightarrow R\text{-mod}_{u\text{-co}}$ preserves direct limits, too.

Now let \mathbf{G} denote the set of all u -comodule left R -modules of the form $\Gamma_u(G)$, where G ranges over (representatives of the isomorphism classes of) all the finitely presented left R -modules. We claim that \mathbf{G} is a set of generators of $R\text{-mod}_{u\text{-co}}$.

Indeed, let M be a u -comodule left R -module; then we have $M \cong \Gamma_u(M)$. Let (G_α) be a diagram of finitely presented left R -modules, indexed by some directed poset, such that $M \cong \varinjlim_\alpha G_\alpha$. Then we have $M \cong \Gamma_u(M) \cong \varinjlim_\alpha \Gamma_u(G_\alpha)$. So M is the direct limit of a diagram of objects from \mathbf{G} in $R\text{-mod}_{u\text{-co}}$, hence it is also a quotient of a coproduct of copies of objects from \mathbf{G} .

The functor Γ_u takes injective objects in $R\text{-mod}$ to injective objects in $R\text{-mod}_{u\text{-co}}$, since it is right adjoint to an exact functor. To show that $\Gamma_u(J)$ is an injective cogenerator of $R\text{-mod}_{u\text{-co}}$ when J is an injective cogenerator of $R\text{-mod}$, it suffices to compute $\text{Hom}_R(M, \Gamma_u(J)) = \text{Hom}_R(M, J) \neq 0$ when $0 \neq M \in R\text{-mod}_{u\text{-co}}$. \square

Lemma 3.5. *Assume that $\text{Tor}_1^R(U, U) = 0$ and $\text{pd}_R U \leq 1$. Then $R\text{-mod}_{u\text{-ctra}}$ is a locally presentable abelian category with a projective generator $\Delta_u(R) \in R\text{-mod}_{u\text{-ctra}}$.*

Proof. Following [22, Example 4.1(1-2)], if λ is a regular cardinal such that the left R -module U is λ -presentable (i. e., isomorphic to the cokernel of a morphism of

free left R -modules with less than λ generators), then the category $R\text{-mod}_{u\text{-ctra}}$ is locally λ -presentable. Since the functor Δ_u is left adjoint to an exact (fully faithful) functor $R\text{-mod}_{u\text{-ctra}} \rightarrow R\text{-mod}$, it takes projective left R -modules to projective u -contramodule left R -modules. Finally, one has $\text{Hom}_R(\Delta_u(R), C) = \text{Hom}_R(R, C) = C \neq 0$ for any object $0 \neq C \in R\text{-mod}_{u\text{-ctra}}$. \square

4. THE ENDOMORPHISM RING OF THE TWO-TERM COMPLEX ($R \rightarrow U$)

According to the discussion in [22, Section 1.1 in the introduction], [23, Section 6.3], and [21, Examples 1.2(4) and 1.3(4)], in the assumptions of Lemma 3.5 the abelian category $\mathbf{B} = R\text{-mod}_{u\text{-ctra}}$ with its natural projective generator $P = \Delta_u(R)$ can be described as the category of modules over an additive monad \mathbb{T}_u on the category of sets. For any set X , the coproduct $P^{(X)}$ of X copies of the object P in the category \mathbf{B} can be computed as $P^{(X)} = \Delta_u(R^{(X)})$, where $R^{(X)} = R[X]$ is the free R -modules with generators indexed by X . The monad \mathbb{T}_u assigns to every set X the set $\text{Hom}_{\mathbf{B}}(P, P^{(X)}) = \Delta_u(R^{(X)})$. In particular, to a one-element set $*$, the monad \mathbb{T}_u assigns the underlying set of the R -module $P = \Delta_u(R)$. In fact $P = \mathbb{T}_u(*) \in \mathbb{T}_u\text{-mod} \cong \mathbf{B}$ is the free \mathbb{T}_u -module with one generator.

For any additive monad \mathbb{T} on the category of sets, the set $\mathbb{T}(*)$ has a natural associative ring structure. This is the ring of endomorphisms of the forgetful functor $\mathbb{T}\text{-mod} \rightarrow \text{Ab}$. In particular, the ring $\mathfrak{R} = \mathbb{T}_u(*)$ can be computed as the opposite ring to the ring of endomorphisms

$$\Delta_u(R) = \text{Ext}_R^1(K^\bullet, R) = \text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, R[1]) \cong \text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, K^\bullet).$$

of the object K^\bullet in the derived category of left R -modules. Notice that the right action of the ring R by endomorphisms of the object $K^\bullet \in \text{D}^b(R\text{-mod})$ in the derived category induces a natural ring homomorphism $R \rightarrow \mathfrak{R}$.

In the next lemma we discuss the particular case of a commutative ring R .

Lemma 4.1. *Let $u: R \rightarrow U$ be an epimorphism of commutative rings such that $\text{Tor}_R^1(U, U) = 0$. Then the ring $\mathfrak{R} = \text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, K^\bullet)$ is commutative. In particular, if u is injective, then the ring $\mathfrak{R} = \text{Hom}_R(U/R, U/R)$ is commutative.*

Proof. This is a generalization of [20, Proposition 3.1]. Let us prove the equivalent assertion that the ring $\mathfrak{R} = \text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet[-1], K^\bullet[-1])$ is commutative (where $K^\bullet[-1]$ is the complex $R \rightarrow U$ with the term R placed in the cohomological degree 0 and the term U placed in the cohomological degree 1). Denote by \mathbf{K} the full subcategory in $\text{D}^b(R\text{-mod})$ consisting of the single object $K^\bullet[-1]$ (and all the objects isomorphic to it). Then the functor of truncated tensor product

$$L^\bullet \bar{\otimes} M^\bullet = \tau_{\geq -1}(L^\bullet \otimes_R^{\mathbb{L}} M^\bullet)$$

defines a unital tensor (monoidal) category structure on the category \mathbf{K} with the unit object $K^\bullet[-1]$. In other words, there is a natural isomorphism $K^\bullet[-1] \bar{\otimes} K^\bullet[-1] \cong$

$K^\bullet[-1]$ transforming both the endomorphisms $f \otimes \text{id}$ and $\text{id} \otimes f$ into the endomorphism f for any $f: K^\bullet[-1] \rightarrow K^\bullet[-1]$. The commutativity of endomorphisms follows formally from that (see the computation in [20]).

When u is a homological epimorphism, one does not need to truncate the tensor product, so one can use the functor $\otimes_R^{\mathbb{L}}$ instead of $\bar{\otimes}$. When u is an injective epimorphism, it suffices to consider the full subcategory spanned by the object $K = U/R$ in $R\text{-mod}$ and the functor $\text{Tor}_1^R(-, -)$ in the role of the tensor product operation. Then one has to use the natural isomorphism $\text{Tor}_1^R(K, K) \cong K$. \square

The next lemma shows that the second assertion of Lemma 4.1 also holds for noninjective ring epimorphisms u of projective dimension ≤ 1 .

Lemma 4.2. *Let $u: R \rightarrow U$ be an epimorphism of associative rings such that $\text{Tor}_R^1(U, U) = 0$ and $\text{pd}_R U \leq 1$. Then the associative ring homomorphism*

$$\text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, K^\bullet) \longrightarrow \text{Hom}_R(U/R, U/R)$$

produced by applying the degree-zero cohomology functor $H^0: \text{D}^b(R\text{-mod}) \rightarrow R\text{-mod}$ to the complex $K^\bullet \in \text{D}^b(R\text{-mod})$ is surjective. In particular, if the ring R is commutative, then so is the ring $\text{Hom}_R(U/R, U/R)$.

Proof. Let $I \subset R$ be the kernel of the map u . Then we have a natural distinguished triangle

$$I[1] \longrightarrow K^\bullet \longrightarrow U/R \longrightarrow I[2]$$

in $\text{D}^b(R\text{-mod-}R)$, and we can also consider it as a distinguished triangle in $\text{D}^b(R\text{-mod})$. Applying the functor $\text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, -[*])$ to this triangle, we see that the map $\text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, K^\bullet) \rightarrow \text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, U/R)$ is surjective, because $\text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, I[2]) = \text{Ext}_R^2(K^\bullet, I) \cong \text{Ext}_R^2(U, I) = 0$ Lemma 2.1(b) and since $\text{pd}_R U \leq 1$. Finally, we have $\text{Hom}_{\text{D}^b(R\text{-mod})}(K^\bullet, U/R) = \text{Ext}_R^0(K^\bullet, U/R) \cong \text{Hom}_R(U/R, U/R)$ by Lemma 2.1(c).

This proves the first assertion of the lemma. The second one follows from the first one together with the first assertion of Lemma 4.1. \square

5. WHEN IS THE CLASS OF TORSION MODULES HEREDITARY?

Notice that every left u -comodule is u -torsion, but the converse implication does not need to be true. The torsion class of all u -torsion left R -modules does *not* need to be hereditary, i. e., a submodule of a u -torsion R -module does not need to be u -torsion. In fact, if $\text{Tor}_1^R(U, U) = 0$ and $\text{fd}_R U \leq 1$, then any one of the mentioned two properties holds if and only if U is a flat right R -module.

Lemma 5.1. *Assume that $\text{Tor}_1^R(U, U) = 0$ and $\text{fd}_R U \leq 1$. Then the following conditions are equivalent:*

- (1) *all u -torsion left R -modules are u -comodules;*
- (2) *all quotient R -modules of left u -comodules are u -comodules;*

- (3) all R -submodules of left u -comodules are u -comodules;
- (4) all R -submodules of u -torsion left R -modules are u -torsion;
- (5) all R -submodules of left u -comodules are u -torsion;
- (6) the right R -module U is flat.

Proof. (1) \implies (2) By the definition, all u -comodules are u -torsion. Hence (1) means that the classes of left u -comodules and u -torsion left R -modules coincide. Since the class of u -torsion R -modules is clearly closed under quotients, (2) follows.

(2) \iff (3) holds because the class of all left u -comodules is closed under kernels and cokernels of morphisms (by Lemma 3.1(a)).

(3) \implies (5) and (4) \implies (5) are obvious.

(5) \implies (6) Let A be a left R -module. From the exact sequence (8) we see that the left R -module $\mathrm{Tor}_1^R(U, A)$ is a submodule of the left R -module $\mathrm{Tor}_1^R(K^\bullet, A)$. By Lemma 2.4(c), $\mathrm{Tor}_1^R(K^\bullet, A) = \Gamma_u(A)$ is a left u -comodule. Under (5), it follows that the left R -module $\mathrm{Tor}_1^R(U, A)$ is u -torsion. Being simultaneously a left U -module, it follows that $\mathrm{Tor}_1^R(U, A) = 0$.

(6) \implies (1) and (6) \implies (4) are obvious. \square

Examples of noncommutative homological ring epimorphisms of projective dimension 1 (on both sides) that are not flat (on either side) do exist. Let k be a field, $k[x]$ be the polynomial ring in one variable x with the coefficients in k , and $kx \subset k[x]$ be the one-dimensional k -vector subspace spanned by x . Then the embedding of matrix rings $R = \begin{pmatrix} k & k \oplus kx \\ 0 & k \end{pmatrix} \longrightarrow \begin{pmatrix} k[x] & k[x] \\ k[x] & k[x] \end{pmatrix} = U$ is an injective ring epimorphism such that $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{pd}_R U = \mathrm{pd} U_R = \mathrm{fd}_R U = \mathrm{fd} U_R = 1$ (cf. Section 8).

On the other hand, the following theorem holds true for epimorphisms of commutative rings.

Theorem 5.2. *If $u: R \longrightarrow U$ is an epimorphism of commutative rings such that $\mathrm{Tor}_1^R(U, U) = 0$ and $\mathrm{pd}_R U \leq 1$, then U is a flat R -module.*

Proof. The argument is based on some results from the papers [11, 1]. Assume first that u is injective. Then $U \oplus U/R$ is a 1-tilting R -module [2, Theorem 3.5], hence $C = \mathrm{Hom}_{\mathbb{Z}}(U \oplus U/R, \mathbb{Q}/\mathbb{Z})$ is a 1-cotilting R -module of cofinite type [9, Theorems 15.2 and 15.18]. The 1-cotilting class associated with C consists of all the R -submodules of U -modules; in other words, it is what we call the class of all u -torsion-free R -modules. Hence the torsion class in the 1-cotilting torsion pair associated with C is the class of all u -torsion R -modules. According to [11, Proposition 3.11], any 1-cotilting torsion pair of cofinite type in the category of modules over a commutative ring is hereditary. By Lemma 5.1 (4) \implies (6), it follows that $\mathrm{fd}_R U = 0$.

In the general case of a (not necessarily injective) homological epimorphism of commutative rings $u: R \longrightarrow U$ with $\mathrm{pd}_R U \leq 1$, one has to use silting theory instead of tilting theory. The R -module $U \oplus U/R$ is 1-silting by [13, Example 6.5], and a 2-term projective resolution of the complex $U \oplus K^\bullet$ is the related silting complex. Hence $C = \mathrm{Hom}_{\mathbb{Z}}(U \oplus U/R, \mathbb{Q}/\mathbb{Z})$ is a cosilting R -module of cofinite type [1, Corollary 3.6]. The cosilting class associated with C consists of all the u -torsion-free R -modules, and

the torsion class in the cosilting torsion pair is the class of all u -torsion R -modules. By [1, Lemma 4.2], any cosilting torsion pair of cofinite type in the category of modules over a commutative ring is hereditary. Once again, by Lemma 5.1 (4) \implies (6) we can conclude that U is a flat R -module. \square

6. TRIANGULATED MATLIS EQUIVALENCE

Let $u: R \rightarrow U$ be a homological epimorphism of associative rings, that is a ring homomorphism such that the natural map of U - U -bimodules $U \otimes_R U \rightarrow U$ is an isomorphism and $\mathrm{Tor}_i^R(U, U) = 0$ for all $i > 0$. Then, according to [8, Theorem 4.4], [16, Theorem 3.7], the restriction of scalars with respect to u is a fully faithful functor between the unbounded derived categories $\mathrm{D}(U\text{-mod}) \rightarrow \mathrm{D}(R\text{-mod})$. We denote this functor, acting between the bounded or unbounded derived categories, by

$$u_*: \mathrm{D}^*(U\text{-mod}) \longrightarrow \mathrm{D}^*(R\text{-mod}),$$

where $\star = \mathbf{b}, +, -, \text{ or } \emptyset$ is a derived category symbol.

In the case of the unbounded derived categories ($\star = \emptyset$), the functor u_* has a left adjoint functor $\mathbb{L}u^*: \mathrm{D}(R\text{-mod}) \rightarrow \mathrm{D}(U\text{-mod})$ and a right adjoint functor $\mathbb{R}u^!: \mathrm{D}(R\text{-mod}) \rightarrow \mathrm{D}(U\text{-mod})$. When U is a right R -module of finite flat dimension, the functor $\mathbb{L}u^*$ also acts between bounded derived categories,

$$\mathbb{L}u^*: \mathrm{D}^*(R\text{-mod}) \longrightarrow \mathrm{D}^*(U\text{-mod}).$$

When U is a left R -module of finite projective dimension, the functor $\mathbb{R}u^!$ acts between bounded derived categories,

$$\mathbb{R}u^!: \mathrm{D}^*(R\text{-mod}) \longrightarrow \mathrm{D}^*(U\text{-mod}).$$

Since the triangulated functor u_* is fully faithful, its left and right adjoints $\mathbb{L}u^*$ and $\mathbb{R}u^!$ are Verdier quotient functors [7, Proposition I.1.3].

Theorem 6.1. (a) *Assume that $\mathrm{fd}U_R \leq 1$. Then the kernel of the functor $\mathbb{L}u^*: \mathrm{D}^*(R\text{-mod}) \rightarrow \mathrm{D}^*(U\text{-mod})$ coincides with the full subcategory $\mathrm{D}_{u\text{-co}}^*(R\text{-mod}) \subset \mathrm{D}^*(R\text{-mod})$ of all complexes of left R -modules with u -comodule cohomology modules. Hence for every symbol $\star = \mathbf{b}, +, -, \text{ or } \emptyset$, we have a triangulated equivalence*

$$\mathrm{D}^*(R\text{-mod})/u_*\mathrm{D}^*(U\text{-mod}) \cong \mathrm{D}_{u\text{-co}}^*(R\text{-mod}).$$

(b) *Assume that $\mathrm{pd}_R U \leq 1$. Then the kernel of the functor $\mathbb{R}u^!: \mathrm{D}^*(R\text{-mod}) \rightarrow \mathrm{D}^*(U\text{-mod})$ coincides with the full subcategory $\mathrm{D}_{u\text{-ctra}}^*(R\text{-mod}) \subset \mathrm{D}^*(R\text{-mod})$ of all complexes of left R -modules with u -contramodule cohomology modules. Hence for every symbol $\star = \mathbf{b}, +, -, \text{ or } \emptyset$, we have a triangulated equivalence*

$$\mathrm{D}^*(R\text{-mod})/u_*\mathrm{D}^*(U\text{-mod}) \cong \mathrm{D}_{u\text{-ctra}}^*(R\text{-mod}).$$

Proof. Part (a): the functor $\mathbb{L}u^*$ is constructed as the derived tensor product $\mathbb{L}u^*(A^\bullet) = U \otimes_R^{\mathbb{L}} A^\bullet$ for any complex of left R -modules A^\bullet . In particular, when $\text{fd } U_R \leq 1$, we have short exact sequences of cohomology

$$0 \longrightarrow U \otimes_R H^n(A^\bullet) \longrightarrow H^n(\mathbb{L}u^*(A^\bullet)) \longrightarrow \text{Tor}_1^R(U, H^{n+1}(A^\bullet)) \longrightarrow 0$$

for any complex $A^\bullet \in \mathbf{D}^*(R\text{-mod})$ and all $n \in \mathbb{Z}$. It follows immediately that $\mathbb{L}u^*(A^\bullet) = 0$ if and only if $H^n(A^\bullet) \in R\text{-mod}_{u\text{-co}}$ for all $n \in \mathbb{Z}$.

Part (b): the functor $\mathbb{R}u^!$ is constructed as the derived homomorphisms $\mathbb{R}u^!(B^\bullet) = \mathbb{R}\text{Hom}_R(U, B^\bullet)$ for any complex of left R -modules B^\bullet . In particular, when $\text{pd } {}_R U \leq 1$, we have short exact sequences of cohomology

$$0 \longrightarrow \text{Ext}_R^1(U, H^{n-1}(B^\bullet)) \longrightarrow H^n(\mathbb{R}u^!(B^\bullet)) \longrightarrow \text{Hom}_R(U, H^n(B^\bullet)) \longrightarrow 0$$

for any complex $B^\bullet \in \mathbf{D}^*(R\text{-mod})$ and all $n \in \mathbb{Z}$. It follows immediately that $\mathbb{R}u^!(B^\bullet) = 0$ if and only if $H^n(B^\bullet) \in R\text{-mod}_{u\text{-ctra}}$ for all $n \in \mathbb{Z}$. \square

Corollary 6.2. *Assume that $\text{fd } U_R \leq 1$ and $\text{pd } {}_R U \leq 1$. Then for every symbol $\star = \mathbf{b}, +, -, \text{ or } \emptyset$ there is a triangulated equivalence*

$$\mathbf{D}_{u\text{-co}}^\star(R\text{-mod}) \cong \mathbf{D}_{u\text{-ctra}}^\star(R\text{-mod})$$

provided by the mutually inverse functors $\mathbb{R}\text{Hom}_R(K^\bullet[-1], -): \mathbf{D}_{u\text{-co}}^\star(R\text{-mod}) \rightarrow \mathbf{D}_{u\text{-ctra}}^\star(R\text{-mod})$ and $K^\bullet[-1] \otimes_R^{\mathbb{L}} -: \mathbf{D}_{u\text{-ctra}}^\star(R\text{-mod}) \rightarrow \mathbf{D}_{u\text{-co}}^\star(R\text{-mod})$.

Proof. More generally, in the context of Theorem 6.1(a), the functor $\mathbf{D}^*(R\text{-mod}) \rightarrow \mathbf{D}_{u\text{-co}}^\star(R\text{-mod})$ right adjoint to the embedding $\mathbf{D}_{u\text{-co}}^\star(R\text{-mod}) \rightarrow \mathbf{D}^*(R\text{-mod})$ is computed as $K^\bullet[-1] \otimes_R^{\mathbb{L}} -$. Similarly, in the context of Theorem 6.1(b), the functor $\mathbf{D}^*(R\text{-mod}) \rightarrow \mathbf{D}_{u\text{-ctra}}^\star(R\text{-mod})$ left adjoint to the embedding $\mathbf{D}_{u\text{-ctra}}^\star(R\text{-mod}) \rightarrow \mathbf{D}^*(R\text{-mod})$ is computed as $\mathbb{R}\text{Hom}_R(K^\bullet[-1], -)$ (cf. [20, Proposition 4.4]). \square

The results of this section can be expressed by existence of the following *recollement* of triangulated categories for any homological ring epimorphism $u: R \rightarrow U$ such that $\text{fd } U_R \leq 1$ and $\text{pd } {}_R U \leq 1$:

$$(10) \quad \begin{array}{c} \begin{array}{ccc} \begin{array}{c} \curvearrowright \\ \leftarrow \\ \curvearrowleft \end{array} & \begin{array}{c} \longrightarrow \\ \longrightarrow \\ \longrightarrow \end{array} & \begin{array}{c} \curvearrowright \\ \leftarrow \\ \curvearrowleft \end{array} \\ \mathbf{D}^*(U\text{-mod}) & \longrightarrow & \mathbf{D}^*(R\text{-mod}) & \longrightarrow & \mathbf{D}_{u\text{-co}}^\star(R\text{-mod}) = \mathbf{D}_{u\text{-ctra}}^\star(R\text{-mod}) \end{array} \end{array}$$

Here the arrows with a tail denote fully faithful triangulated functors, while the arrows with two heads denote triangulated Verdier quotient functors. The two leftmost curvilinear arrows are adjoint on the left and on the right to the leftmost straight arrow, while the two rightmost curvilinear arrows are adjoint on the left and on the right to the rightmost straight arrow. The image of the leftmost straight arrow is the kernel of the rightmost straight arrow, and similarly with the two pairs of curvilinear arrows.

The three leftmost arrows are the functors $\mathbb{L}u^*$, u_* , and $\mathbb{R}u^!$. The two rightmost curvilinear arrows are the inclusions of the full subcategories $\mathbf{D}_{u\text{-co}}^\star(R\text{-mod})$ and $\mathbf{D}_{u\text{-ctra}}^\star(R\text{-mod})$ into $\mathbf{D}^*(R\text{-mod})$.

7. TWO FULLY FAITHFUL TRIANGULATED FUNCTORS

In addition to the assumptions on the projective and flat dimension of the left and right R -module U that we used above, the results of this section require certain assumptions about the properties of injective and projective left R -modules vis-à-vis the homological ring homomorphism $u: R \rightarrow U$. Specifically, these are the assumptions that injective left R -modules are u -special and projective left R -modules are u -cospecial, or in other words, the left R -modules $\mathrm{Tor}_0^R(K^\bullet, J) = U/R \otimes_R J$ and $\mathrm{Ext}_R^0(K^\bullet, F) = \mathrm{Hom}_R(U/R, F)$ vanish for all injective left R -modules J and projective left R -modules F (cf. Lemmas 2.1(c) and 2.2).

Theorem 7.1. (a) *Assume that $\mathrm{fd} U_R \leq 1$ and $(U/R) \otimes_R J = 0$ for all injective left R -modules J . Then, for any conventional derived category symbol $\star = \mathbf{b}, +, -, \text{ or } \emptyset$, the triangulated functor*

$$D^\star(R\text{-mod}_{u\text{-co}}) \longrightarrow D^\star(R\text{-mod})$$

induced by the exact embedding of abelian categories $R\text{-mod}_{u\text{-co}} \rightarrow R\text{-mod}$ is fully faithful, and its essential image coincides with the full subcategory

$$D_{u\text{-co}}^\star(R\text{-mod}) \subset D^\star(R\text{-mod}),$$

providing an equivalence of triangulated categories

$$D^\star(R\text{-mod}_{u\text{-co}}) \cong D_{u\text{-co}}^\star(R\text{-mod}).$$

(b) *Assume that $\mathrm{pd}_R U \leq 1$ and $\mathrm{Hom}_R(U/R, F) = 0$ for all projective left R -modules F . Then, for any conventional derived category symbol $\star = \mathbf{b}, +, -, \text{ or } \emptyset$, the triangulated functor*

$$D^\star(R\text{-mod}_{u\text{-ctra}}) \longrightarrow D^\star(R\text{-mod})$$

induced by the exact embedding of abelian categories $R\text{-mod}_{u\text{-ctra}} \rightarrow R\text{-mod}$ is fully faithful, and its essential image coincides with the full subcategory

$$D_{u\text{-ctra}}^\star(R\text{-mod}) \subset D^\star(R\text{-mod}),$$

providing an equivalence of triangulated categories

$$D^\star(R\text{-mod}_{u\text{-ctra}}) \cong D_{u\text{-ctra}}^\star(R\text{-mod}).$$

Proof. This is an application of the general technique formulated in [20, Theorem 6.4 and Proposition 6.5]. Let us explain part (b). The pair of functors $\mathrm{Ext}_R^i(K^\bullet, -)$, $i = 0, 1$, is a cohomological functor between the abelian categories $R\text{-mod}$ and $R\text{-mod}_{u\text{-ctra}}$, that is, for every short exact sequence of left R -modules $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ there is a short exact sequence of left u -contramodules (cf. Lemmas 2.1(a-b) and 2.5(c))

$$\begin{aligned} 0 \longrightarrow \mathrm{Ext}_R^0(K^\bullet, A) \longrightarrow \mathrm{Ext}_R^0(K^\bullet, B) \longrightarrow \mathrm{Ext}_R^0(K^\bullet, C) \\ \longrightarrow \mathrm{Ext}_R^1(K^\bullet, A) \longrightarrow \mathrm{Ext}_R^1(K^\bullet, B) \longrightarrow \mathrm{Ext}_R^1(K^\bullet, C) \longrightarrow 0. \end{aligned}$$

Since, by our assumption, the functor $\text{Ext}_R^0(K^\bullet, -)$ annihilates projective left R -modules, it follows that our cohomological functor $\text{Ext}_R^*(K^\bullet, -)$ is the left derived functor of the functor $\Delta = \Delta_u = \text{Ext}_R^1(K^\bullet, -): R\text{-mod} \rightarrow R\text{-mod}_{u\text{-ctra}}$, that is $\mathbb{L}_1\Delta_u = \text{Ext}_R^0(K^\bullet, -)$ and $\mathbb{L}_i\Delta_u = 0$ for $i > 1$.

By Lemma 3.2(b), the functor Δ_u is left adjoint to the exact, fully faithful embedding functor $R\text{-mod}_{u\text{-ctra}} \rightarrow R\text{-mod}$, so we are in the setting of [20, Theorem 6.4]. It remains to point out that $\mathbb{L}_1\Delta_u(B) = \text{Ext}_R^0(K^\bullet, B) = 0$ for all left u -contramodules B . Notice that the class $R\text{-mod}_{\Delta\text{-adj}} = \text{Ker}(\mathbb{L}_{>0}\Delta)$ of Δ -adjusted left R -modules, playing a key role in the argument in [20, Section 6], is nothing but the class of u -cospecial left R -modules in our context, according to Lemma 2.2.

Similarly, in part (a) one observes that the pair of functors $\text{Tor}_i^R(K^\bullet, -)$, $i = 0, 1$ is a homological functor between the abelian categories $R\text{-mod}$ and $R\text{-mod}_{u\text{-co}}$, hence, whenever the functor $\text{Tor}_1^R(K^\bullet, -)$ annihilates injective left R -modules, it is the right derived functor of the functor $\Gamma = \Gamma_u = \text{Tor}_1^R(K^\bullet, -): R\text{-mod} \rightarrow R\text{-mod}_{u\text{-co}}$, that is $\mathbb{R}^1\Gamma_u = \text{Tor}_0^R(K^\bullet, -)$ and $\mathbb{R}^i\Gamma = 0$ for $i > 1$. It remains to point out that $\mathbb{R}^1\Gamma_u(A) = \text{Tor}_0^R(K^\bullet, A) = 0$ for all left u -comodules A . As above, we notice that the class $R\text{-mod}_{\Gamma\text{-adj}} = \text{Ker}(\mathbb{R}^{>0}\Gamma)$ of Γ -adjusted left R -modules is just the class of u -special left R -modules discussed in Section 2. \square

Remark 7.2. Conversely, if $\text{fd } U_R \leq 1$ and the triangulated functor $\text{D}^b(R\text{-mod}_{u\text{-co}}) \rightarrow \text{D}^b(R\text{-mod})$ is fully faithful, then $(U/R) \otimes_R J = 0$ for all injective left R -modules J . A proof of this can be found in [5, Lemma 3.9 and Proposition 4.2] (cf. [20, Remark 6.8]). Similarly, if $\text{pd}_R U \leq 1$ and the triangulated functor $\text{D}^b(R\text{-mod}_{u\text{-ctra}}) \rightarrow \text{D}^b(R\text{-mod})$ is fully faithful, then $\text{Hom}_R(U/R, F) = 0$ for all projective left R -modules F [5, Lemma 3.9 and Proposition 4.1].

The following result can be also found in [5, Corollary 4.4].

Corollary 7.3. *Let $u: R \rightarrow U$ be a homological ring epimorphism. Assume that $\text{fd } U_R \leq 1$ and $\text{pd}_R U \leq 1$. Suppose further that $(U/R) \otimes_R J = 0$ for all injective left R -modules J and $\text{Hom}_R(U/R, F) = 0$ for all projective left R -modules F . Then for every conventional derived category symbol $\star = \mathbf{b}, +, -, \text{ or } \emptyset$, there is a triangulated equivalence between the derived categories of the abelian categories $R\text{-mod}_{u\text{-co}}$ and $R\text{-mod}_{u\text{-ctra}}$ of left u -comodules and left u -contramodules,*

$$\text{D}^\star(R\text{-mod}_{u\text{-co}}) \cong \text{D}^\star(R\text{-mod}_{u\text{-ctra}}).$$

Proof. According to Corollary 6.2 and Theorem 7.1(a-b), we have a chain of triangulated equivalences

$$\text{D}^\star(R\text{-mod}_{u\text{-co}}) \cong \text{D}_{u\text{-co}}^\star(R\text{-mod}) \cong \text{D}_{u\text{-ctra}}^\star(R\text{-mod}) \cong \text{D}^\star(R\text{-mod}_{u\text{-ctra}}).$$

\square

In the assumptions of Corollary 7.3, the recollement (10) takes the form

$$(11) \quad \begin{array}{c} \begin{array}{ccc} \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} \\ \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} \\ \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} \end{array} \\ \begin{array}{ccc} \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} \\ \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} \\ \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} & \begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \end{array} \end{array} \end{array}$$

In the recollement (11), all the three triangulated categories are derived categories of certain abelian categories (and the third one is even the derived category of two different abelian categories).

Example 7.4. For any injective ring epimorphism $u: R \rightarrow U$, the conditions $(U/R) \otimes_R J = 0$ and $\text{Hom}_R(U/R, F) = 0$ hold for all injective left R -modules J and all projective left R -modules F . Indeed, if u is injective and J is an injective left R -module, then any left R -module morphism $R \rightarrow J$ can be extended to a left R -module morphism $U \rightarrow J$. Hence the left R -module J is u -h-divisible (i. e., a quotient R -module of a left U -module). Thus $U/R \otimes_R U = 0$ implies $U/R \otimes_R J = 0$. Similarly, the map $F \rightarrow U \otimes_R F$ is injective for any flat left R -module F , so F is u -torsion-free (i. e., an R -submodule of a left U -module). Therefore, $\text{Hom}_R(U/R, U) = 0$ implies $\text{Hom}_R(U/R, F) = 0$.

8. KRONECKER QUIVER EXAMPLE

Let k be an algebraically closed field, and let R denote the path algebra of the Kronecker quiver $\bullet \rightrightarrows \bullet$ over k . So left R -modules are pairs of k -vector spaces (V_1, V_2) endowed with a pair of k -linear maps $f_V, g_V: V_1 \rightrightarrows V_2$. The aim of this section is to describe the full subcategories of comodules and contramodules for certain ring epimorphisms originating from R .

We will interpret R as the matrix ring $R = \begin{pmatrix} k & k \oplus kx \\ 0 & k \end{pmatrix}$, where the element $1 \in k \oplus kx$ in the upper right corner acts in the quiver representations by the map f_V and the element $x \in k \oplus kx$ acts by the map g_V . When the map f_V is invertible, the fraction $x = g_V/f_V$ is a linear operator $V_1 \rightarrow V_1$ or $V_2 \rightarrow V_2$. The eigenvalues of this operator, if they happen to exist, can be thought of as points of the projective line $\mathbb{P}^1(k) = k \cup \{\infty\}$ with the coordinate x .

Let $\mathbb{X} \subset \mathbb{P}^1(k)$ be a subset of points of the projective line such that $\infty \in \mathbb{X}$. Denote by $S_{\mathbb{X}} = \mathbb{X}^{-1}k[x]$ the localization of the ring of polynomials $k[x]$ at the multiplicative subset generated by the elements $x - \lambda$, $\lambda \in \mathbb{X} \setminus \{\infty\}$. Consider the matrix ring $U_{\mathbb{X}} = \begin{pmatrix} S_{\mathbb{X}} & S_{\mathbb{X}} \\ S_{\mathbb{X}} & S_{\mathbb{X}} \end{pmatrix}$. Then there is a ring homomorphism $u_{\mathbb{X}}: R \rightarrow U_{\mathbb{X}}$ given by the inclusion of the matrices. The map $u_{\mathbb{X}}$ is a homological ring epimorphism. The essential image of the functor of restriction of scalars $u_{\mathbb{X}*}: U_{\mathbb{X}}\text{-mod} \rightarrow R\text{-mod}$ consists of all the quiver representations (f_V, g_V) such that the map f_V is an isomorphism and the map $g_V - \lambda f_V: V_1 \rightarrow V_2$ is an isomorphism for all $\lambda \in \mathbb{X} \setminus \{\infty\}$.

In particular, for $\mathbb{X} = \{\infty\}$ we have $U_{\{\infty\}} = \begin{pmatrix} k[x] & k[x] \\ k[x] & k[x] \end{pmatrix}$. The essential image of the functor $u_{\{\infty\}*}: U_{\{\infty\}}\text{-mod} \rightarrow R\text{-mod}$ consists of all the quiver representations

(f_V, g_V) such that the map f_V is invertible. For an arbitrary subset $\{\infty\} \in \mathbb{X} \subset \mathbb{P}^1(k)$, the morphism $u_{\mathbb{X}}: R \rightarrow U_{\mathbb{X}}$ factorizes as the composition of two injective homological ring epimorphisms $R \rightarrow U_{\{\infty\}} \rightarrow U_{\mathbb{X}}$. The ring $U_{\mathbb{X}}$ has both flat and projective dimension 1 both as a left and as a right R -module (as a left and right $U_{\{\infty\}}$ -module, it has flat dimension 0 and projective dimension 1).

So the full categories of $u_{\mathbb{X}}$ -comodules and $u_{\mathbb{X}}$ -contramodules in $R\text{-mod}$ are abelian. Specifically, let us say that a quiver representation $M = (f_M, g_M)$ is a “ $(\lambda = \infty)$ -comodule” if $\text{Hom}_R(M, V) = 0 = \text{Ext}_R^1(M, V)$ for any quiver representation $V = (f_V, g_V)$ with the map f_V invertible. A quiver representation $C = (f_C, g_C)$ is a “ $(\lambda = \infty)$ -contramodule” if $\text{Hom}_R(V, C) = 0 = \text{Ext}_R^1(V, C)$ for any such V .

More generally, we will say that a quiver representation M is an “ \mathbb{X} -comodule” if $\text{Hom}_R(M, V) = 0 = \text{Ext}_R^1(M, V)$ for any $V = (f_V, g_V)$ with the maps f_V and $g_V - \lambda f_V$ invertible for all $\lambda \in \mathbb{X}$. A quiver representation C is an “ \mathbb{X} -contramodule” if $\text{Hom}_R(V, C) = 0 = \text{Ext}_R^1(V, C)$ for any such V . One can easily see that a quiver representation is an \mathbb{X} -comodule if and only if the related left R -module is a $u_{\mathbb{X}}$ -comodule, and similarly, a quiver representation is an \mathbb{X} -contramodule if and only if the related left R -module is a $u_{\mathbb{X}}$ -contramodule.

Assume first that $0 \notin \mathbb{X}$. Denote by y the coordinate $1/x$ on $\mathbb{P}^1(k)$ (so y is a possible eigenvalue of f_V/g_V), and let $\mathbb{Y} \subset \mathbb{A}^1(k)$ denote the subset of the affine line consisting of all points $\mu \in k$ such that $\mu^{-1} \in \mathbb{X}$. Consider the polynomial ring $k[y]$, and denote by $T_{\mathbb{Y}} = \mathbb{Y}^{-1}k[y]$ its localization at the multiplicative subset generated by the elements $y - \mu$, $\mu \in \mathbb{Y}$. Denote by $v_{\mathbb{Y}}$ the ring epimorphism $k[y] \rightarrow T_{\mathbb{Y}}$. In particular, if $\mathbb{X} = \{\infty\}$, then $\mathbb{Y} = \{0\}$ and $T_{\mathbb{Y}} = k[y, y^{-1}]$.

The results of the following lemma are (easy) particular cases of the theory developed in [19, Section 13] and [3, Section 4 and/or 6].

Lemma 8.1. (a) *If $\mathbb{Y} = \{\mu\}$ is a one-point set, then a $k[y]$ -module M is a $v_{\{\mu\}}$ -comodule if and only if the operator $y - \mu$ is locally nilpotent in M , i. e., for every $m \in M$ there exists $n \in \mathbb{Z}_{\geq 1}$ such that $(y - \mu)^n m = 0$. For an arbitrary subset $\mathbb{Y} \subset \mathbb{A}^1(k)$, any $v_{\mathbb{Y}}$ -comodule M has a unique, functorial decomposition into a direct sum of $v_{\{\mu\}}$ -comodules M_{μ} over $\mu \in \mathbb{Y}$, and any such direct sum $M = \bigoplus_{\mu \in \mathbb{Y}} M_{\mu}$ of $v_{\{\mu\}}$ -comodules M_{μ} is a $v_{\mathbb{Y}}$ -comodule. The category $k[y]\text{-mod}_{v_{\mathbb{Y}}\text{-co}}$ of $v_{\mathbb{Y}}$ -comodules is thus equivalent to the Cartesian product of the categories $k[y]\text{-mod}_{v_{\{\mu\}}\text{-co}}$ over $\mu \in \mathbb{Y}$.*

(b) *If $\mathbb{Y} = \{\mu\}$ is a one-point set, then a $k[y]$ -module C is a $v_{\{\mu\}}$ -contramodule if and only if it admits $(y - \mu)$ -power infinite summation operations in the sense of [19, Section 3]. For an arbitrary subset $\mathbb{Y} \subset \mathbb{A}^1(k)$, any $v_{\mathbb{Y}}$ -contramodule C has a unique, functorial decomposition into a direct product of $v_{\{\mu\}}$ -contramodules C^{μ} over $\mu \in \mathbb{Y}$, and any such direct product $C = \prod_{\mu \in \mathbb{Y}} C^{\mu}$ of $v_{\{\mu\}}$ -contramodules C^{μ} is a $v_{\mathbb{Y}}$ -contramodule. The category $k[y]\text{-mod}_{v_{\mathbb{Y}}\text{-contra}}$ of $v_{\mathbb{Y}}$ -contramodules is thus equivalent to the Cartesian product of the categories $k[y]\text{-mod}_{v_{\{\mu\}}\text{-contra}}$ over $\mu \in \mathbb{Y}$. \square*

The next proposition describes \mathbb{X} -comodule and \mathbb{X} -contramodule quiver representations for $\mathbb{X} \subset \mathbb{P}^1(k)$, $\infty \in \mathbb{X}$, $0 \notin \mathbb{X}$.

Proposition 8.2. *Let \mathbb{X} be a subset in $\mathbb{P}^1(k) = k \cup \{\infty\}$ containing ∞ and not containing 0, and let $\mathbb{Y} \subset \mathbb{A}_k^1 = k$ be the set of all μ such that $\mu^{-1} \in \mathbb{X}$. Then*

(a) *a Kronecker quiver representation $M = (f_M, g_M)$ is an \mathbb{X} -comodule if and only if the map g_M is invertible and the vector space $M_1 \cong M_2$ with the linear operator $y = f_M/g_M$ is a $v_{\mathbb{Y}}$ -comodule;*

(b) *a Kronecker quiver representation $C = (f_C, g_C)$ is an \mathbb{X} -contramodule if and only if the map g_C is invertible and the vector space $C_1 \cong C_2$ with the linear operator $y = f_C/g_C$ is a $v_{\mathbb{Y}}$ -contramodule.*

It follows from Lemma 8.1 and Proposition 8.2 that the category of \mathbb{X} -comodules decomposes as a Cartesian product of \mathbb{X} copies of the category of vector spaces with a locally nilpotent operator z , and similarly, the category of \mathbb{X} -contramodules decomposes as a Cartesian product of \mathbb{X} copies of the category of vector spaces with z -power infinite summation operations.

Notice that it follows from Proposition 8.2 that the category of \mathbb{X} -comodules is *equivalent* to a torsion class (viewed as a full subcategory) in $k[y]\text{-mod}$. But a subrepresentation of an \mathbb{X} -comodule is *not* an \mathbb{X} -comodule, generally speaking (because the condition of invertibility of the operator g_M is not preserved by the passage to a subrepresentation), in agreement with the discussion in Section 5.

The proof of Proposition 8.2 given below consists of several lemmas.

Lemma 8.3. *For any \mathbb{X} not containing 0, one has:*

(a) *in any \mathbb{X} -comodule $M = (f_M, g_M)$, the map g_M is invertible;*

(b) *in any \mathbb{X} -contramodule $C = (f_C, g_C)$, the map g_C is invertible.*

Proof. We will prove part (b). Assume that the operator g_C has a nonzero kernel $K_1 \subset C_1$. Then there are two possibilities. If the kernel of the restriction of f_C to K_1 is nonzero, then C contains a copy of the injective representation $k \rightrightarrows 0$ as a subrepresentation. In this case, for any nonzero representation V with f_V invertible (hence $V_1 \neq 0$) there exists a nonzero morphism $V \rightarrow (k \rightrightarrows 0) \rightarrow C$. If the map $f_C|_{K_1}$ is injective, then C contains a nonzero subrepresentation $K = (K_1, f_C(K_1))$ with f_K invertible and $g_K = 0$, hence $g_K - \lambda f_K$ is invertible for all $\lambda \in \mathbb{X}$. In both cases, there exists a $U_{\mathbb{X}}$ -module V and a nonzero morphism $V \rightarrow C$, contradicting the assumption that C is an \mathbb{X} -contramodule.

Assume that the map g_C is not surjective. Then the quiver representation C has a quotient representation $L = (C_1, C_2/g_C(C_2))$ with $g_L = 0$ and $L_2 = \text{coker}(g_C) \neq 0$. Once again, there are two possibilities. If the map f_L is not surjective, then C has a projective quotient representation $N = (0 \rightrightarrows k)$. In this case, for any quiver representation V , one has $\text{Ext}_R^1(V, N) = 0$ if and only if V is projective. In particular, $\text{Ext}_R^1(V, N) \neq 0$ for any nonzero representation V with f_V invertible.

If the map f_L is surjective, then $N = (L_1/\ker(f_L), L_2)$ is a nonzero quotient representation of C with $g_N = 0$ and f_N invertible. In this case, it suffices to notice that a nonzero vector space with a zero operator $x = g/f$ is not an injective object of the category of $k[x]$ -modules. In particular, consider the quiver representation $V = (k \rightrightarrows k)$ with $f_V = 1$ and $g_V = 0$ (so f_V is invertible and $g_V - \lambda f_V$ is invertible

for all $\lambda \in \mathbb{X}$). Then $\text{Ext}_R^1(V, N) = \text{Ext}_{k[x]}^1(k, N_1) \cong N_1 \neq 0$ (where $x = g/f$ acts by zero both in k and in N_1).

In both cases, we have found a $U_{\mathbb{X}}$ -module V such that $\text{Ext}_R^1(V, N) \neq 0$. Since the category of Kronecker quiver representations has homological dimension 1, the functor $\text{Ext}_R^1(V, -)$ is right exact and it follows that $\text{Ext}_R^1(V, C) \neq 0$. \square

Lemma 8.4. *For any \mathbb{X} not containing 0, one has:*

(a) *for any \mathbb{X} -comodule $M = (f_M, g_M)$, the vector space $M_1 \cong M_2$ with the linear operator $y = f_M/g_M$ is a $v_{\mathbb{Y}}$ -comodule;*

(b) *for any \mathbb{X} -contramodule $C = (f_C, g_C)$, the vector space $C_1 \cong C_2$ with the linear operator $y = f_C/g_C$ is a $v_{\mathbb{Y}}$ -contramodule.*

Proof. Part (b): For any quiver representations V and C with the operators g_V and g_C invertible one has $\text{Ext}_R^*(V, C) \cong \text{Ext}_{k[y]}^*(V_1, C_1)$, where y acts in $V_1 \cong V_2$ and $C_1 \cong C_2$ by the operators f/g . Set $V_1 = T_{\mathbb{Y}} = V_2$, with the operator f_V being the multiplication with y and $g_V = \text{id}$. Then the maps f_V and g_V are invertible, and so is the map $g_V - \lambda f_V$ for all $\lambda \in \mathbb{X}$. Hence $\text{Ext}_{k[y]}^*(T_{\mathbb{Y}}, C_1) \cong \text{Ext}_R^*(V, C) = 0$ whenever C is an \mathbb{X} -contramodule. The proof of part (a) is similar. \square

Lemma 8.5. *For any \mathbb{X} not containing 0, one has:*

(a) *any Kronecker quiver representation $M = (f_M, g_M)$ such that the map g_M is invertible and the vector space $M_1 \cong M_2$ with the operator $y = f_M/g_M$ belongs to $k[y]\text{-mod}_{v_{\mathbb{Y}\text{-co}}}$ is an \mathbb{X} -comodule;*

(b) *any Kronecker quiver representation $C = (f_C, g_C)$ such that the map g_C is invertible and the vector space $C_1 \cong C_2$ with the operator $y = f_C/g_C$ belongs to $k[y]\text{-mod}_{v_{\mathbb{Y}\text{-ctra}}}$ is an \mathbb{X} -contramodule.*

Proof. Part (b): the class of \mathbb{X} -contramodules is closed under infinite products in the category of Kronecker quiver representations. Hence, in view of Lemma 8.1(b), it suffices to consider the case when the vector space $C_1 \cong C_2$ with the operator $y = f_C/g_C$ belongs to $k[y]\text{-mod}_{v_{\{\mu\}\text{-ctra}}}$ for some fixed value of $\mu \in \mathbb{Y}$. Changing the coordinate on $\mathbb{P}^1(k)$ reduces the question to the case $\mu = 0$.

Furthermore, any $v_{\{0\}}$ -contramodule (or in other words, a k -vector space with a y -power infinite summation operation) can be obtained from the 1-dimensional vector space k with the operator $y = 0$ using cokernels, extensions, and projective limits (of which the latter reduce to kernels and infinite products). The class of \mathbb{X} -contramodules is closed under all those operations in the category of quiver representations; so it suffices to show that the representations C with g_C invertible and $f_C = 0$ are \mathbb{X} -contramodules for all $\mathbb{X} \ni \infty$. Without loss of generality, one can assume that $\mathbb{X} = \{\infty\}$.

Let V be a Kronecker quiver representation with f_V invertible. We have to check that $\text{Hom}_R(V, C) = 0 = \text{Ext}_R^1(V, C)$. Indeed, any morphism $h: V \rightarrow C$ vanishes, since the invertibility of f_V and the vanishing of f_C together imply the vanishing of the map $h_2: V_2 \rightarrow C_2$, and then in view of the invertibility of g_C the map $h_1: V_1 \rightarrow C_1$ vanishes as well. Now let $0 \rightarrow C \rightarrow A \rightarrow V \rightarrow 0$ be a short exact sequence of quiver representations. Then the subspaces C_2 and $f_A(A_1)$ form a

direct sum decomposition of the vector space A_2 , and it follows that the subspaces C_1 and $g_A^{-1}(f_A(A_1))$ form a direct sum decomposition of the vector space A_1 . Thus the short exact sequence of representations is split.

The proof of part (a) is similar. \square

Proof of Proposition 8.2. Follows from Lemmas 8.3, 8.4, and 8.5. \square

Now we consider the general case when the subset $\mathbb{X} \subset \mathbb{P}^1(k)$ may contain the point 0 (so one can possibly have $\mathbb{X} = \mathbb{P}^1(k)$). The idea is to compute the R - R -bimodule $K = U_{\mathbb{X}}/R$, and consequently the functors $\Gamma_{u_{\mathbb{X}}} = \text{Tor}_1^R(K, -)$ and $\Delta_{u_{\mathbb{X}}} = \text{Ext}_R^1(K, -)$. Then we will use the following category-theoretic observations.

Lemma 8.6. *Let \mathcal{C} be a category with products and a zero object. Assume that the identity endofunctor $\text{Id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$ decomposes as a product of a family of functors $F_i: \mathcal{C} \rightarrow \mathcal{C}$, where i ranges over some index set I ,*

$$\text{Id}_{\mathcal{C}} = \prod_{i \in I} F_i.$$

Denote by $\mathcal{C}_i \subset \mathcal{C}$ the essential image of the functor F_i , viewed as a full subcategory in \mathcal{C} . Then the category \mathcal{C} is equivalent to the Cartesian product of the categories \mathcal{C}_i ,

$$\mathcal{C} \cong \prod_{i \in I} \mathcal{C}_i.$$

Proof. The functor $F: \mathcal{C} \rightarrow \prod_{i \in I} \mathcal{C}_i$ is simply the collection of the functors F_i , that is $F = (F_i)_{i \in I}$. The inverse functor $G: \prod_{i \in I} \mathcal{C}_i \rightarrow \mathcal{C}$ is the functor of I -indexed product in the category \mathcal{C} restricted to the full subcategory $\prod_{i \in I} \mathcal{C}_i \subset \mathcal{C}^I$. Clearly, the composition $G \circ F: \mathcal{C} \rightarrow \mathcal{C}$ is the identity functor.

The key observation is that for any objects $C, D \in \mathcal{C}$, any morphism $\prod_{i \in I} F_i(C) \cong C \rightarrow D \cong \prod_{i \in I} F_i(D)$ decomposes as a product of morphisms $F_i(C) \rightarrow F_i(D)$. It follows that, for any objects $C_i \in \mathcal{C}_i$ and $D_j \in \mathcal{C}_j$ with $i \neq j$, there are no nonzero morphisms $C_i \rightarrow D_j$. Hence for any object $C_i \in \mathcal{C}_i$ one has $F_j(C_i) = 0$ for all $j \neq i$, and consequently $F_i(C_i) = C_i$. Furthermore, the functors F_i preserve products in \mathcal{C} , since they are retracts of the identity functor. This allows to show that the composition $F \circ G$ is the identity functor. \square

We recall that, in the category-theoretic terminology, a left adjoint functor to the inclusion of a subcategory is called a *reflector*, and a subcategory admitting such an adjoint functor is said to be *reflective*.

Lemma 8.7. *Let \mathcal{A} be a category with products and a zero object, and let $\mathcal{C} \subset \mathcal{A}$ be a reflective full subcategory with the reflector $\Delta: \mathcal{A} \rightarrow \mathcal{C}$. Suppose that the functor Δ decomposes as a product of a family of functors $\Delta_i: \mathcal{A} \rightarrow \mathcal{C}$,*

$$\Delta = \prod_{i \in I} \Delta_i.$$

Denote by $\mathcal{C}_i \subset \mathcal{A}$ the essential image of the functor Δ_i , viewed as a full subcategory in \mathcal{A} . Then one has $\mathcal{C}_i \subset \mathcal{C}$, the category \mathcal{C} is equivalent to the Cartesian product of the categories \mathcal{C}_i , and the functor $\Delta_i: \mathcal{A} \rightarrow \mathcal{C}_i$ is the reflector onto the full subcategory $\mathcal{C}_i \subset \mathcal{A}$.

Proof. Set $F_i = \Delta_i|_C$ and apply the previous lemma. \square

The following theorem is the main result of this section.

Theorem 8.8. *For any subset $\infty \in \mathbb{X} \subset \mathbb{P}^1(k)$, the following assertions hold.*

(a) *Any \mathbb{X} -comodule M has a unique, functorial decomposition into a direct sum of $\{\lambda\}$ -comodules M_λ over $\lambda \in \mathbb{X}$, and any such direct sum $M = \bigoplus_{\lambda \in \mathbb{X}} M_\lambda$ of $\{\lambda\}$ -comodules M_λ is a \mathbb{X} -comodule. The category of \mathbb{X} -comodules is thus equivalent to the Cartesian product of the categories of $\{\lambda\}$ -comodules over $\lambda \in \mathbb{X}$ (each of which is equivalent to the category of k -vector spaces with a locally nilpotent linear operator z).*

(b) *Any \mathbb{X} -contramodule C has a unique, functorial decomposition into a direct product of $\{\lambda\}$ -contramodules C^λ over $\lambda \in \mathbb{X}$, and any such direct product $C = \prod_{\lambda \in \mathbb{X}} C^\lambda$ of $\{\lambda\}$ -contramodules C^λ is a \mathbb{X} -contramodule. The category of \mathbb{X} -contramodules is thus equivalent to the Cartesian product of the categories of $\{\lambda\}$ -contramodules over $\lambda \in \mathbb{X}$ (each of which is equivalent to the category of k -vector spaces with a z -power infinite summation operation).*

Proof. The R - R -bimodule $U_{\mathbb{X}}$ can be described as the following representation of the Cartesian square $(\bullet \rightrightarrows \bullet) \times (\bullet \rightrightarrows \bullet)$ of the Kronecker quiver $\bullet \rightrightarrows \bullet$ (we recall the notation $S_{\mathbb{X}} = \mathbb{X}^{-1}k[x]$ for the relevant localization of the polynomial ring $k[x]$):

$$(12) \quad \begin{array}{ccc} S_{\mathbb{X}} & \begin{array}{c} \xrightarrow{1} \\ \xrightarrow{x} \end{array} & S_{\mathbb{X}} \\ \begin{array}{c} \uparrow 1 \\ \uparrow x \end{array} & & \begin{array}{c} \uparrow 1 \\ \uparrow x \end{array} \\ S_{\mathbb{X}} & \begin{array}{c} \xrightarrow{1} \\ \xrightarrow{x} \end{array} & S_{\mathbb{X}} \end{array}$$

In the same vein, the R - R -bimodule $K = U_{\mathbb{X}}/R$ is described as the following representation of the quiver $(\bullet \rightrightarrows \bullet) \times (\bullet \rightrightarrows \bullet)$:

$$(13) \quad \begin{array}{ccc} S_{\mathbb{X}}/k & \begin{array}{c} \xrightarrow{1} \\ \xrightarrow{x} \end{array} & S_{\mathbb{X}}/(k \oplus kx) \\ \begin{array}{c} \uparrow 1 \\ \uparrow x \end{array} & & \begin{array}{c} \uparrow 1 \\ \uparrow x \end{array} \\ S_{\mathbb{X}} & \begin{array}{c} \xrightarrow{1} \\ \xrightarrow{x} \end{array} & S_{\mathbb{X}}/k \end{array}$$

The key observation is that the representation (13) decomposes into a direct sum of representations indexed by the points of the set \mathbb{X} . This direct sum decomposition of (13) is induced by the direct sum decomposition

$$S_{\mathbb{X}} \cong k[x] \oplus \bigoplus_{\lambda \in \mathbb{X} \setminus \{\infty\}} \left(\bigoplus_{n \geq 1} k(x - \lambda)^{-n} \right)$$

of the vector space $S_{\mathbb{X}}$. Hence we obtain an \mathbb{X} -indexed direct sum decomposition of the functor $\Gamma_{u_{\mathbb{X}}} = \text{Tor}_1^R(K, -)$ and an \mathbb{X} -indexed direct product decomposition of the

functor $\Delta_{u_{\mathbb{X}}} = \text{Ext}_R^1(K, -)$. It remains to apply Proposition 3.1(b) together with the dual assertion to Lemma 8.7 in order to deduce part (a) of the theorem, and Proposition 3.2(b) together with Lemma 8.7 in order to deduce part (b).

The description of the categories of $\{\lambda\}$ -comodules and $\{\lambda\}$ -contramodules in parts (a) and (b) is provided by Proposition 8.2. It suffices to change the coordinate on the projective line $\mathbb{P}^1(k)$ suitably in order to include the case $\lambda = 0$. \square

REFERENCES

- [1] L. Angeleri Hügel, M. Hrbek. Silting modules over commutative rings. *Internat. Math. Research Notices* **2017**, #13, p. 4131–4151. [arXiv:1602.04321](#) [math.RT]
- [2] L. Angeleri Hügel, J. Sánchez. Tilting modules arising from ring epimorphisms. *Algebras and Represent. Theory* **14**, #2, p. 217–246, 2011. [arXiv:0804.1313](#) [math.RT]
- [3] S. Bazzoni, L. Positselski. S -almost perfect commutative rings. *Journ. of Algebra* **532**, p. 323–356, 2019. [arXiv:1801.04820](#) [math.AC]
- [4] S. Bazzoni, J. Šťovíček. Smashing localizations of rings of weak global dimension at most one. *Advances in Math.* **305**, p. 351–401, 2017. [arXiv:1402.7294](#) [math.AC]
- [5] H.-X. Chen, C.-C. Xi. Derived decompositions of abelian categories I. Electronic preprint [arXiv:1804.10759](#) [math.RT].
- [6] A. Facchini, Z. Nazemian. Equivalence of some homological conditions for ring epimorphisms. *Journ. of Pure and Appl. Algebra* **223**, #4, p. 1440–1455, 2019. [arXiv:1710.00097](#) [math.RA]
- [7] P. Gabriel, M. Zisman. Calculus of fractions and homotopy theory. Springer-Verlag, Berlin–Heidelberg–New York, 1967.
- [8] W. Geigle, H. Lenzing. Perpendicular categories with applications to representations and sheaves. *Journ. of Algebra* **144**, #2, p. 273–343, 1991.
- [9] R. Göbel, J. Trlifaj. Approximations and endomorphism algebras of modules. Second revised and extended edition. De Gruyter, 2012.
- [10] D. K. Harrison. Infinite abelian groups and homological methods. *Annals of Math.* **69**, #2, p. 366–391, 1959.
- [11] M. Hrbek. One-tilting classes and modules over commutative rings. *Journ. of Algebra* **462**, p. 1–22, 2016. [arXiv:1507.02811](#) [math.AC]
- [12] H. Krause, J. Šťovíček. The telescope conjecture for hereditary rings via Ext-orthogonal pairs. *Advances in Math.* **225**, #5, p. 2341–2364, 2010. [arXiv:0810.1401](#) [math.RT]
- [13] F. Marks, J. Šťovíček. Universal localizations via silting. *Proc. of the Royal Soc. of Edinburgh, Sect. A* **149**, #2, p. 511–532, 2019. [arXiv:1605.04222](#) [math.RT]
- [14] E. Matlis. Cotorsion modules. *Memoirs of the American Math. Society* **49**, 1964.
- [15] E. Matlis. 1-dimensional Cohen–Macaulay rings. Lecture notes in Math., 327, Springer-Verlag, Berlin–Heidelberg–New York, 1973.
- [16] D. Pauksztello. Homological epimorphisms of differential graded algebras. *Communicat. in Algebra* **37**, #7, p. 2337–2350, 2009. [arXiv:0705.0118](#) [math.RA]
- [17] L. Positselski. Contramodules. Electronic preprint [arXiv:1503.00991](#) [math.CT].
- [18] L. Positselski. Dedualizing complexes and MGM duality. *Journ. of Pure and Appl. Algebra* **220**, #12, p. 3866–3909, 2016. [arXiv:1503.05523](#) [math.CT]
- [19] L. Positselski. Contraadjusted modules, contramodules, and reduced cotorsion modules. *Moscow Math. Journ.* **17**, #3, p. 385–455, 2017. [arXiv:1605.03934](#) [math.CT]
- [20] L. Positselski. Triangulated Matlis equivalence. *Journ. of Algebra and its Appl.* **17**, #4, article ID 1850067, 2018. [arXiv:1605.08018](#) [math.CT]

- [21] L. Positselski. Abelian right perpendicular subcategories in module categories. Electronic preprint [arXiv:1705.04960](https://arxiv.org/abs/1705.04960) [math.CT].
- [22] L. Positselski, J. Rosický. Covers, envelopes, and cotorsion theories in locally presentable abelian categories and contramodule categories. *Journ. of Algebra* **483**, p. 83–128, 2017. [arXiv:1512.08119](https://arxiv.org/abs/1512.08119) [math.CT]
- [23] L. Positselski, J. Šťovíček. The tilting-cotilting correspondence. *Internat. Math. Research Notices*, published online at <https://doi.org/10.1093/imrn/rnz116> in July 2019. [arXiv:1710.02230](https://arxiv.org/abs/1710.02230) [math.CT]
- [24] B. Stenström. Rings of quotients. An introduction to methods of ring theory. Springer-Verlag, Berlin–Heidelberg–New York, 1975.

(Silvana Bazzoni) DIPARTIMENTO DI MATEMATICA “TULLIO LEVI-CIVITA”, UNIVERSITÀ DI PADOVA, VIA TRIESTE 63, 35121 PADOVA (ITALY)

E-mail address: bazzoni@math.unipd.it

(Leonid Positselski) INSTITUTE OF MATHEMATICS OF THE CZECH ACADEMY OF SCIENCES, ŽITNÁ 25, 115 67 PRAHA 1 (CZECH REPUBLIC); AND

LABORATORY OF ALGEBRA AND NUMBER THEORY, INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS, MOSCOW 127051 (RUSSIA)

E-mail address: positselski@yandex.ru