

Lab exercises

Prof. S. De Marchi

Padova, May 8, 2018

1 Recalls about trapezoidal and Simpson's formulas

For the approximation of definite integrals, we saw two important methods: the *trapezoid formula* and the *Simpson's formula*. Both are *interpolatory* formulas of Newton-Côtes type.

We have studied their *simple* expressions and the *composite or generalized* ones.

Let $I(f) = \int_a^b f(x)dx$ be the integral we want to compute.

Letting $I_T(f)$, $I_T^c(f)$ and $I_S(f)$, $I_S^c(f)$ the corresponding *simple* and *composite* formulas, we have

T. **Simple trapezoid formula and its error.**

$$I_T(f) = \frac{b-a}{2} (f(a) + f(b)) , \quad (1)$$

$$E_T(f) = I(f) - I_T(f) = -\frac{(b-a)^3}{12} f''(\xi) , \quad \xi \in (a, b).$$

Tc. **Trapezoidal formula and its error.**

Here we must take an equispaced subdivision of $[a, b]$ such as $\{x_0 = a, \dots, x_i = a + ih, \dots, x_n = b\}$, with $h = (b-a)/n$:

$$I_T^c(f) = \frac{h}{2} (f(a) + f(b)) + h \sum_{i=1}^{n-1} f(x_i) , \quad (2)$$

$$E_T^c(f) = I(f) - I_T^c(f) = -\frac{(b-a)^3}{12n^2} f''(\xi) , \quad \xi \in (a, b).$$

S. **Simple Simpson's formula and its error.**

$$I_S(f) = \frac{b-a}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] , \quad (3)$$

$$E_S(f) = I(f) - I_S(f) = -\frac{1}{24} \frac{(b-a)^5}{32} f^{(4)}(\xi) , \quad \xi \in (a, b) , \quad h = (b-a)/2 .$$

Sc. **Composite Simpson's formula and its error.**

Here we take, as in the case of the trapezoids formula, an equispaced subdivision of $[a, b]$. Hence in the general interval $I_k = [x_{k-1}, x_k]$, we consider the points x_{k-1} , $\bar{x}_k = \frac{x_{k-1}+x_k}{2}$

and x_k . Setting $h = (b - a)/(2n)$, this is equivalent to take the points $2n + 1$ points $a = z_0, z_1, z_2, \dots, z_{2n} = b$.

we have:

$$I_S^c(f) = \frac{h}{3}(f(z_0) + f(z_{2n})) + \frac{h}{3} \left[4 \sum_{k=1}^{2n-1''} f(z_k) + 2 \sum_{k=2}^{2n-2''} f(z_k) \right], \quad (4)$$

where the " indicate that the sum is done either on the odd or the even indeces, respectively

$$E_S^c(f) = I(f) - I_S^c(f) = -\frac{(b-a)^5}{2880n^4} f^{(4)}(\xi), \quad \xi \in (a, b).$$

2 Proposed exercises

1. Write the M-function that implements the trapezoidal formula (2). **Notice** that for $n = 1$ it gives the formula (1).
2. Write the M-function that implements the composite Simpson formula (4). Again **notice** that for $n = 1$ it gives the formula (3).
3. Compute the approximation of

$$\int_{-1}^1 \frac{x}{2} e^{-\frac{x}{2}} \cos(x) dx \quad (5)$$

by means of the simple Simpson's formula and calculate the absolute error. As exact value, take the output of the Matlab function

```
quadl(inline('x/2.*exp(-x/2).*cos(x)'), -1, 1, 1.e-9)
```

4. Compute numerically

$$\int_0^{2\pi} x e^{-x} \cos(2x) dx = \frac{3(e^{-2\pi} - 1) - 10\pi e^{-2\pi}}{25} \approx -0.12212260462,$$

by means of **composite** formulas of trapezoids and Simpson,

- (a) Find *a priori* the number of quadrature points so that the absolute errors $E_T^c(f)$ and $E_S^c(f)$ be in modulus less than `tol = 1.e - 5`. (*Hint*: find the max absolute value of $f^{(2)}$ and $f^{(4)}$ by using a fine discretization of the interval $[0, 2\pi]$)
- (b) Determine also the absolute error with respect to the exact value of the integral.

Time: **2 hours**.