

The generalized recurrent set, explosions and Lyapunov functions

OLGA BERNARDI¹ ANNA FLORIO² JIM WISEMAN³

¹ Dipartimento di Matematica “Tullio Levi-Civita”, Università di Padova, Italy

² Laboratoire de Mathématiques d’Avignon, Avignon Université, France

³ Department of Mathematics, Agnes Scott College, Decatur, Georgia, USA

Abstract

We consider explosions in the generalized recurrent set for homeomorphisms on a compact metric space. We provide multiple examples to show that such explosions can occur, in contrast to the case for the chain recurrent set. We give sufficient conditions to avoid explosions and discuss their necessity. Moreover, we explain the relations between explosions and cycles for the generalized recurrent set. In particular, for a compact topological manifold with dimension greater or equal 2, we characterize explosion phenomena in terms of existence of cycles. We apply our results to give sufficient conditions for stability, under \mathcal{C}^0 perturbations, of the property of admitting a continuous Lyapunov function which is not a first integral.

1 Introduction

Generalized recurrence was originally introduced for flows by Auslander in the Sixties [3] by using continuous Lyapunov functions. Auslander defined the generalized recurrent set to be the union of those orbits along which all continuous Lyapunov functions are constant. In the same paper, he gave a characterization of this set in terms of the theory of prolongations. The generalized recurrent set was later extended to maps by Akin and Auslander (see [1] and [2]). More recently Fathi and Pageault [7] proved that, for a homeomorphism, the generalized recurrent set can be equivalently defined by using Easton’s strong chain recurrence [6].

The present paper is concerned with the behaviour, under continuous perturbations of the map, of the generalized recurrent set for homeomorphisms. In particular, we analyze the phenomenon of explosions, which are discontinuous jumps in the size of the generalized recurrent set. Moreover, we apply our results to give sufficient conditions to assure the persistence under continuous perturbations of a continuous Lyapunov function which is not a first integral (that is a continuous strict Lyapunov function).

Throughout the paper, (X, d) is a compact metric space. We denote by $Hom(X)$ the space of homeomorphisms of X equipped with the uniform topology induced by the metric:

$$d_{\mathcal{C}^0}(f, g) = \max_{x \in X} d(f(x), g(x)).$$

Let $\mathcal{GR}(f)$ be the generalized recurrent set of $f \in Hom(X)$ (the rigorous definition will be given in Section 2).

Assume that $\mathcal{GR}(f) \neq X$. We say f does not permit \mathcal{GR} -explosions if for any open neighborhood U of $\mathcal{GR}(f)$ in X there exists a neighborhood V of f in $Hom(X)$ such that if $g \in V$ then $\mathcal{GR}(g) \subset U$.

We say f does not permit \mathcal{GR} -full explosions if there exists a neighborhood V of f in $Hom(X)$ such that if $g \in V$ then $\mathcal{GR}(g) \neq X$.

Explosions have been studied for both the non-wandering set $\mathcal{NW}(f)$ and the chain recurrent set $\mathcal{CR}(f)$

(see e.g. [15] and [4] respectively). We introduce the notion of \mathcal{GR} -full explosion for the application to continuous strict Lyapunov functions. After some preliminary results and definitions, in Section 2 we observe that \mathcal{GR} -(full) explosions in general can occur (Example 2.1).

Section 3 gives sufficient conditions to avoid \mathcal{GR} -explosions (Corollary 3.1 and Proposition 3.2) and \mathcal{GR} -full explosions (Proposition 3.3). These results can be summarized as follows.

Assume that $\mathcal{GR}(f) \neq X$. If (i) $\mathcal{GR}(f) = \mathcal{CR}(f)$ or (ii) f is topologically stable then f does not permit \mathcal{GR} -explosions. If $\mathcal{CR}(f) \neq X$ then f does not permit \mathcal{GR} -full explosions.

The core of Section 3 concerns the necessity of these conditions. In particular, in Examples 3.2, 3.3 and 3.4 respectively, we show that the converses of the previous results are in general false on compact metric spaces.

The goal of Section 4 is to explain the relations between explosions and cycles for the generalized recurrent set. The notion of cycle will be rigorously recalled in Definition 4.1. Since we need to apply the \mathcal{C}^0 closing lemma, the ambient space is a compact, topological manifold M with $\dim(M) \geq 2$.

Assume that $\mathcal{GR}(f) \neq M$. f does not permit \mathcal{GR} -explosions if and only if there exists a decomposition of $\mathcal{GR}(f)$ with no cycles.

The above theorem generalizes the corresponding result for the non-wandering set. More precisely, the fact that the existence of a decomposition of $\mathcal{NW}(f)$ without cycles prohibits \mathcal{NW} -explosions is due to Pugh and Shub (see Theorem 6.1 in [14] for flows and Theorem 5.6 in [18] for homeomorphisms); the converse has been proved by Shub and Smale (see Lemma 2 in [15]). We postpone the proofs of both implications to Appendixes A and B.

We remark that, in the proof of this theorem, we do not use the full definition of the generalized recurrent set, but only the fact that $\mathcal{GR}(f)$ is a compact, invariant set, which contains $\mathcal{NW}(f)$ and is contained in $\mathcal{CR}(f)$. Consequently, with the same proof, we obtain the corresponding results both for the strong chain recurrent set $\mathcal{SCR}_d(f)$ and $\mathcal{CR}(f)$, see Theorem 4.5 and Remark 4.2.

In Section 5 we apply the results of Sections 3 and 4 to obtain sufficient conditions for an affirmative answer to the question:

For a given homeomorphism, is the property of admitting a continuous strict Lyapunov function stable under \mathcal{C}^0 perturbations?

We refer to Propositions 5.1 and 5.2. Finally, in Proposition 5.3, we remark that on a smooth, compact manifold the property of admitting such a function is generic.

2 Preliminaries

In this section we recall the notions of chain recurrent, strong chain recurrent and generalized recurrent point for a fixed $f \in \text{Hom}(X)$.

Given $x, y \in X$ and $\varepsilon > 0$, an ε -chain from x to y is a finite sequence $(x_i)_{i=1}^n \subset X$ such that $x_1 = x$ and, setting, $x_{n+1} = y$, we have

$$d(f(x_i), x_{i+1}) < \varepsilon \quad \forall i = 1, \dots, n. \quad (1)$$

A point $x \in X$ is said to be *chain recurrent* if for all $\varepsilon > 0$ there exists an ε -chain from x to x . The set of chain recurrent points is denoted by $\mathcal{CR}(f)$. Since we assumed X to be compact, chain recurrence depends only on the topology, not on the choice of the metric (see for example [5][Theorem 4.4.5] and [8][Section 1]).

Given $x, y \in X$ and $\varepsilon > 0$, a *strong ε -chain* from x to y is a finite sequence $(x_i)_{i=1}^n \subset X$ such that $x_1 = x$ and, setting $x_{n+1} = y$, we have

$$\sum_{i=1}^n d(f(x_i), x_{i+1}) < \varepsilon. \quad (2)$$

A point $x \in X$ is said to be *strong chain recurrent* if for all $\varepsilon > 0$ there exists a strong ε -chain from x to x . The set of strong chain recurrent points is denoted by $\mathcal{SCR}_d(f)$. In general, strong chain recurrence depends on the choice of the metric; see for example [21][Example 3.1] and [19][Example 2.6].

A way to eliminate the dependence on the metric in $\mathcal{SCR}_d(f)$ is taking the intersection over all metrics. We then obtain the generalized recurrent set

$$\mathcal{GR}(f) := \bigcap_d \mathcal{SCR}_d(f) \quad (3)$$

where the intersection is over all metrics compatible with the topology of X .

The sets $\mathcal{GR}(f)$, $\mathcal{SCR}_d(f)$ and $\mathcal{CR}(f)$ are all closed and invariant (see respectively [2][Page 52], [7][Page 1193] and [8][Section 1]). Moreover, in general $\mathcal{NW}(f) \subset \mathcal{GR}(f) \subset \mathcal{SCR}_d(f) \subset \mathcal{CR}(f)$, where $\mathcal{NW}(f)$ denotes the non-wandering set of f , and all inclusions can be strict. We refer to [19][Example 2.9] for an exhaustive treatment of these inclusions.

The dynamical relevance of the generalized recurrent set relies on its relations with continuous Lyapunov functions. A continuous function $u : X \rightarrow \mathbb{R}$ is a Lyapunov function for f if $u(f(x)) \leq u(x)$ for every $x \in X$. Given a Lyapunov function $u : X \rightarrow \mathbb{R}$ for f , the corresponding neutral set is given by

$$\mathcal{N}(u) = \{x \in X : u(f(x)) = u(x)\}.$$

We refer to [7][Theorem 3.1] for the proof of the next result.

Theorem 2.1.

$$\mathcal{GR}(f) = \bigcap_{u \in \mathcal{L}(f)} \mathcal{N}(u),$$

where $\mathcal{L}(f)$ is the set of continuous Lyapunov functions for f . Moreover, there exists a continuous Lyapunov function for f such that $\mathcal{N}(u) = \mathcal{GR}(f)$.

In order to describe the behavior of $\mathcal{GR}(f)$ under continuous perturbations of f , we introduce and discuss the notions of \mathcal{GR} -explosion and \mathcal{GR} -full explosion.

We start by recalling the phenomenon of explosions for the generalized recurrent set, which are particular discontinuities of the function

$$\text{Hom}(X) \ni f \mapsto \mathcal{GR}(f) \in \mathcal{P}(X),$$

where $\mathcal{P}(X)$ denotes the power set of X .

Definition 2.1. (No \mathcal{GR} -explosions) Let $f \in \text{Hom}(X)$ be such that $\mathcal{GR}(f) \neq X$. We say f does not permit \mathcal{GR} -explosions if for any open neighborhood U of $\mathcal{GR}(f)$ in X there exists a neighborhood V of f in $\text{Hom}(X)$ such that if $g \in V$ then $\mathcal{GR}(g) \subset U$.

No \mathcal{NW} -explosions and no \mathcal{CR} -explosions are defined analogously (see for example [15][Page 588] for $\mathcal{NW}(f)$ and [4][Page 323] for $\mathcal{CR}(f)$). We recall that \mathcal{NW} -explosions in general can occur; see [13], [11][Section 6.3] and [18][Section 5.2]. This is not the case for \mathcal{CR} -explosions; see the following, which is Theorem F in [4].

Theorem 2.2. Let $f \in \text{Hom}(X)$ be such that $\mathcal{CR}(f) \neq X$. Then f does not permit \mathcal{CR} -explosions.

In [4], the proof of the previous result essentially uses a dynamical characterization of the points outside $\mathcal{CR}(f)$. We propose an alternative, direct proof of this fact.

Proof of Theorem 2.2. Argue by contradiction and suppose there are an open neighborhood U of $\mathcal{CR}(f)$ in X , a sequence $(g_n)_{n \in \mathbb{N}} \in \text{Hom}(X)$ converging to f in the uniform topology and a sequence of points $(y_n)_{n \in \mathbb{N}}$ such that $y_n \in \mathcal{CR}(g_n) \setminus U$. Since X is compact and U is open, we can assume that the sequence $(y_n)_{n \in \mathbb{N}}$ converges to $y \notin U$. In particular, y does not belong to $\mathcal{CR}(f)$.

Let $\varepsilon > 0$ be fixed. By hypothesis, for any $n \in \mathbb{N}$ there exists an $\frac{\varepsilon}{3}$ -chain $(x_1 = y_n, x_2, \dots, x_m, x_{m+1} = y_n)$ for g_n from y_n to y_n . Corresponding to $\varepsilon > 0$, there exists $\bar{n} \in \mathbb{N}$ such that

$$d_{\mathcal{C}^0}(f, g_n) < \frac{\varepsilon}{3} \quad \forall n \geq \bar{n}. \quad (4)$$

Moreover, by the uniform continuity of f , there exists $\delta \in (0, \frac{\varepsilon}{3})$ such that

$$d(z, w) < \delta \quad \Rightarrow \quad d(f(z), f(w)) < \frac{\varepsilon}{3}. \quad (5)$$

Finally, let $\tilde{n} \in \mathbb{N}$ such that

$$d(y, y_n) < \delta \quad \forall n \geq \tilde{n}. \quad (6)$$

We will show that, if $n \geq \max(\bar{n}, \tilde{n})$ then the chain

$$(y, x_2, \dots, x_m, y)$$

is an ε -chain for f from y to y . Indeed, thanks to inequalities (4),(5) and (6) and the fact that $(x_1 = y_n, x_2, \dots, x_m, x_{m+1} = y_n)$ is an $\frac{\varepsilon}{3}$ -chain for g_n , we have

$$d(f(y), x_2) \leq d(f(y), f(y_n)) + d(f(y_n), g_n(y_n)) + d(g_n(y_n), x_2) < \varepsilon.$$

Moreover

$$d(f(x_i), x_{i-1}) \leq d(f(x_i), g_n(x_i)) + d(g_n(x_i), x_{i+1}) < \frac{2\varepsilon}{3} < \varepsilon$$

for all $i = 2, \dots, m-1$. Finally

$$d(f(x_m), y) \leq d(f(x_m), g_n(x_m)) + d(g_n(x_m), y_n) + d(y_n, y) < \varepsilon.$$

By the arbitrariness of $\varepsilon > 0$, we conclude that $y \in \mathcal{CR}(f)$, obtaining the desired contradiction. \square

We now introduce full explosions for the generalized recurrent set.

Definition 2.2. (No \mathcal{GR} -full explosions) Let $f \in \text{Hom}(X)$ be such that $\mathcal{GR}(f) \neq X$. We say f does not permit \mathcal{GR} -full explosions if there exists a neighborhood V of f in $\text{Hom}(X)$ such that if $g \in V$ then $\mathcal{GR}(g) \neq X$.

Clearly, if f does not permit \mathcal{GR} -explosions then f does not permit \mathcal{GR} -full explosions; if f permits \mathcal{GR} -full explosions then f permits \mathcal{GR} -explosions.

We observe that, unlike the chain recurrent case, \mathcal{GR} -(full) explosions can in general occur.

Example 2.1. On the circle \mathbb{S}^1 with the usual topology, consider an interval $I \subsetneq \mathbb{S}^1$ of positive Lebesgue measure. Let $\phi : \mathbb{S}^1 \rightarrow [0, +\infty)$ be a non-negative smooth function whose set of zeroes is I . Let $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ be the time-one map of the flow of the vector field

$$V(x) = \phi(x) \frac{\partial}{\partial x}.$$

(See Figure 1). In such a case

$$I = \mathcal{GR}(f).$$

We observe that, for an arbitrarily small $\varepsilon > 0$, there always exists $g_\varepsilon \in \text{Hom}(\mathbb{S}^1)$ such that

$$d_{\mathcal{C}^0}(f, g_\varepsilon) < \varepsilon$$

and

$$\mathbb{S}^1 = \mathcal{GR}(g_\varepsilon).$$

(Simply perturb ϕ so that it is positive on I). Consequently, $f \in \text{Hom}(\mathbb{S}^1)$ is an example of a homeomorphism which permits \mathcal{GR} -full explosions. In particular, f admits \mathcal{GR} -explosions.

3 \mathcal{GR} -(full) explosions

The aim of this section is to discuss some sufficient conditions to avoid \mathcal{GR} -(full) explosions.

The first one comes from a straightforward application of Theorem 2.2 (see Corollary G in [4] for $\mathcal{NW}(f)$).

Corollary 3.1. Let $f \in \text{Hom}(X)$ be such that $\mathcal{GR}(f) \neq X$. If $\mathcal{GR}(f) = \mathcal{CR}(f)$ then f does not permit \mathcal{GR} -explosions.

Notice that the converse of Corollary 3.1 is false, in general. In the next two examples, we define homeomorphisms which do not permit \mathcal{GR} -explosions even though $\mathcal{GR}(f) \neq \mathcal{CR}(f)$.




Figure 1: The dynamical system of Example 2.1.

Example 3.1. Let \mathbb{S}^1 be the circle with the usual topology. We consider it as the interval $I = [0, 1]$ with the endpoints identified. Let $K \subset \mathbb{S}^1$ be the middle-third Cantor set constructed on the interval I . Denote as $\{e_n\}_{n \in \mathbb{N}} \subset K$ the set of endpoints of the removed intervals. At each e_n , glue n copies of the interval I . Let X be the union of \mathbb{S}^1 with all these attached copies of I . Define the homeomorphism $f : X \rightarrow X$ as follows: f fixes K and every copy of the interval I , f moves all the points in $\mathbb{S}^1 \setminus K$ counterclockwise (see Figure 2). Clearly, it holds that

$$\mathcal{CR}(f) = X.$$

Moreover, since there is a Cantor set of fixed points,

$$\mathcal{GR}(f) = \text{Fix}(f).$$

We refer to [7][Example 3.3] for details on this argument. The idea is that, even if K has vanishing Lebesgue measure, the dynamical system is topologically conjugate to the case $\lambda_{\text{Leb}}(K) > 0$, in which case strong chains cannot cross K .

We observe that every $g \in \text{Hom}(X)$ must fix K . Indeed, since the e_n 's have homeomorphically distinct neighborhoods, any homeomorphism g must fix each e_n . Moreover, since the endpoints are dense in K , g must fix the entire Cantor set. As a consequence, the homeomorphism f does not permit \mathcal{GR} -explosions.




Figure 2: The dynamical system of Example 3.1.

Example 3.2. We construct the compact metric space X as follows. Let \mathbb{S}^1 be the circle with the usual topology. For notational convenience, we consider it as the interval $[-2, 2]$ with the endpoints identified. Let $K \subset \mathbb{S}^1$ be the middle-third Cantor set constructed on the interval $[-1, 1]$. Denote the set $K \cap [0, 1]$ by K_+ and the set $K \cap [-1, 0]$ by K_- . Observe that the element of K_+ closest to 0 is $1/3$ and the element

of K_- closest to 0 is $-1/3$. For each point $k \in K_+$, let $X_{-k,k}$ be an arc (disjoint from \mathbb{S}^1) connecting $-k$ in K_- to $k \in K_+$. Assume that these arcs are pairwise disjoint. Finally, let

$$X = \mathbb{S}^1 \cup \left(\bigcup_{k \in K_+} X_{-k,k} \right).$$

Define the homeomorphism $f : X \rightarrow X$ as follows. On \mathbb{S}^1 , f fixes K and moves all other points counterclockwise, that is, in the direction of decreasing x . On every arc $X_{-k,k}$, the endpoints $-k$ and k are fixed, since they are in K , and f moves all other points from $-k \in K_-$ toward $k \in K_+$. See Figure 3. In such a case,

$$\mathcal{CR}(f) = X.$$

Moreover, since there is a Cantor set of fixed points,

$$\mathcal{GR}(f) = K \cup (-1/3, 1/3) \cup X_{-1/3, 1/3}.$$

We claim that f does not permit \mathcal{GR} -explosions. To see this, let g be a homeomorphism close to f . Then $g(K) = K$ and each $X_{-k,k}$ maps to some $X_{-k',k'}$. Consequently, if g moves $k \in K_+$ clockwise then g moves $-k \in K_-$ counterclockwise; if g moves $k \in K_+$ counterclockwise then g moves $-k \in K_-$ clockwise. For any $\alpha > 0$, we can assume g close enough to f that

- (a) g moves counterclockwise any $x \in \mathbb{S}^1$ that is not within α of K ;
- (b) For any $x \in X_{-k,k}$ that is not within α of K , $g(x)$ is closer to $k \in K_+$ than is x .

We will show that any $x \in X$ not within α of $\mathcal{GR}(f)$ is not generalized recurrent. Since α is arbitrary, this means that f does not permit \mathcal{GR} -explosions.

For arbitrary $a, b \in \mathbb{S}^1$, we indicate by $[a, b]$ the closed interval in \mathbb{S}^1 obtained by connecting clockwise a and b . Let x be a point not within α of $\mathcal{GR}(f)$. Then the following three cases can occur:

- (i) The point $x \in [0, 2]$ (the right-side semicircle in Figure 3). Then, recalling that points along the arcs $X_{-k,k}$ move from K_- toward K_+ , any chain from x back to itself must go counterclockwise to $[-2, 0]$ (the left-side semicircle in Figure 3). If every point of $K_+ \cap [0, x]$ is fixed, then for some metric and some $\varepsilon > 0$ there is no strong ε -chain from x to $[-2, 0]$, and thus x cannot be generalized recurrent. Otherwise, there must be a point $k' \in K_+ \cap [0, x]$ that is not fixed. If g moves k' clockwise, then the interval $[k', x]$ maps into its interior. As a consequence, for small enough $\varepsilon > 0$ no ε -chain can get from $[k', x]$ to the left-side semicircle, and so x is not chain recurrent. In particular, x is not generalized recurrent. If g moves k' counterclockwise, then g moves $-k'$ clockwise. Then the interval $[-k', x]$ maps into its interior, and again x is not chain recurrent and therefore not generalized recurrent.
- (ii) The point $x \in [-2, 0]$. To return to itself, it would first have to pass through $[0, 2]$. Then the argument in (i) shows that x cannot return to $[-2, 0]$, and thus x is not generalized recurrent.
- (iii) The point $x \in X_{-k,k}$ for some $k \in K_+$. Again, since points along the arcs $X_{-k,k}$ move from K_- toward K_+ , the only way for the point x to come back to itself is passing through $[0, 2]$. Then the argument in (i) shows that x cannot belong to $\mathcal{GR}(g)$.

Summarizing, $f \in \text{Hom}(X)$ is an example of a homeomorphism such that $\mathcal{GR}(f) \neq \mathcal{CR}(f)$ and f does not permit \mathcal{GR} -explosions.

The other implication is valid on a compact topological manifold M with $\dim(M) \geq 2$; see Proposition 3.1 below (see also Theorem H in [4] for $\mathcal{NW}(f)$). To prove this fact we need the following \mathcal{C}^0 closing lemma; see [4][Lemmas 4 and 5] and [12][Lemma 13].

Lemma 3.1. *Let $f \in \text{Hom}(M)$ be defined on a compact topological manifold M of $\dim(M) \geq 2$. If $x \in \mathcal{CR}(f)$ then for any neighborhood V of f in $\text{Hom}(M)$ there exists $g \in V$ such that $x \in \text{Per}(g)$.*

Proposition 3.1. *Let $f \in \text{Hom}(M)$ be defined on a compact topological manifold M of $\dim(M) \geq 2$ and such that $\mathcal{GR}(f) \neq M$. $\mathcal{GR}(f) = \mathcal{CR}(f)$ if and only if f does not permit \mathcal{GR} -explosions.*




Figure 3: The dynamical system of Example 3.2.

Proof. One direction is exactly Corollary 3.1. Assume now that f does not permit \mathcal{GR} -explosions, which means that for any open neighborhood U of $\mathcal{GR}(f)$ there exists a neighborhood V of f in $\text{Hom}(M)$ such that for any $g \in V$ we have $\mathcal{GR}(g) \subset U$. Then we have:

$$\mathcal{GR}(f) \subseteq \mathcal{CR}(f) \subseteq \bigcup_{g \in V} \text{Per}(g) \subseteq \bigcup_{g \in V} \mathcal{GR}(g) \subseteq U,$$

where the second inclusion comes from the \mathcal{C}^0 closing lemma, here Lemma 3.1. Since the neighborhood U of $\mathcal{GR}(f)$ is arbitrary, we conclude that $\mathcal{GR}(f) = \mathcal{CR}(f)$. \square

The second sufficient condition which avoids \mathcal{GR} -explosions is topological stability. We recall the notion of topologically stable homeomorphism, see e.g. [17][Definition 5].

Definition 3.1. A homeomorphism $f \in \text{Hom}(X)$ is topologically stable if there exists a neighborhood V of f in $\text{Hom}(X)$ such that for each $g \in V$ there is a continuous function $h_g : X \rightarrow X$ satisfying:

- (i) $h_g \circ g = f \circ h_g$ (semi-conjugation);
- (ii) $h_g \rightarrow \text{id}$ as $g \rightarrow f$ in the uniform topology.

Proposition 3.2. Let $f \in \text{Hom}(X)$ be such that $\mathcal{GR}(f) \neq X$. If f is topologically stable then f does not permit \mathcal{GR} -explosions.

In order to prove Proposition 3.2, we need the next result.

Lemma 3.2. Let $f, g \in \text{Hom}(X)$. If there exists a continuous function $h : X \rightarrow X$ such that

$$h \circ g = f \circ h$$

then $h(\mathcal{GR}(g)) \subseteq \mathcal{GR}(f)$.

Proof. Recall that, by Theorem 2.1,

$$\mathcal{GR}(f) = \bigcap_{u \in \mathcal{L}(f)} \mathcal{N}(u)$$

where the intersection is taken over the set $\mathcal{L}(f)$ of continuous Lyapunov functions for f .

Let us show that if $u \in \mathcal{L}(f)$ then $u \circ h \in \mathcal{L}(g)$. Indeed for any $z \in X$, since $h \circ g = f \circ h$ and $u \in \mathcal{L}(f)$, we have

$$u \circ h \circ g(z) = u \circ f \circ h(z) \leq u \circ h(z).$$

Now let $z = h(x) \in h(\mathcal{GR}(g))$ for some $x \in \mathcal{GR}(g)$ and consider $u \in \mathcal{L}(f)$. Since $h \circ g = f \circ h$, $x \in \mathcal{GR}(g)$ and –as remarked above– $u \circ h \in \mathcal{L}(g)$, we deduce that

$$u \circ f(z) = u \circ f \circ h(x) = u \circ h \circ g(x) = u \circ h(x) = u(z).$$

This means that $z = h(x) \in \mathcal{N}(u)$. By the arbitrariness of $u \in \mathcal{L}(f)$ and by Theorem 2.1, we conclude that $z \in \mathcal{GR}(f)$. Equivalently, $h(\mathcal{GR}(g)) \subseteq \mathcal{GR}(f)$. \square

We now prove Proposition 3.2.

Proof of Proposition 3.2. Argue by contradiction and suppose there are an open neighborhood U of $\mathcal{GR}(f)$ in X , a sequence $(g_n)_{n \in \mathbb{N}} \in \text{Hom}(X)$ converging to f in the uniform topology and a sequence of points $(x_n)_{n \in \mathbb{N}}$ such that $x_n \in \mathcal{GR}(g_n) \setminus U$.

Since f is topologically stable and the sequence $(g_n)_{n \in \mathbb{N}}$ converges uniformly to f , there exists an index $\bar{n} \in \mathbb{N}$ such that

$$h_n \circ g_n = f \circ h_n \quad \forall n \geq \bar{n}$$

where $h_n : X \rightarrow X$ is a continuous map. Consequently, by Lemma 3.2,

$$h_n(\mathcal{GR}(g_n)) \subseteq \mathcal{GR}(f) \quad \forall n \geq \bar{n}. \quad (7)$$

Moreover, by Definition 3.1, the sequence $(h_n)_{n \in \mathbb{N}}$ converges uniformly to id . Define the following continuous function

$$X \setminus U \ni y \mapsto d(y, \mathcal{GR}(f)) := \min_{x \in \mathcal{GR}(f)} d(y, x).$$

Since $X \setminus U$ is compact and $(X \setminus U) \cap \mathcal{GR}(f) = \emptyset$, it holds that $\eta := \min_{y \in X \setminus U} d(y, \mathcal{GR}(f)) > 0$. Let now $\tilde{n} \in \mathbb{N}$, $\tilde{n} \geq \bar{n}$ be such that

$$d_{\mathcal{G}^0}(id, h_n) < \eta \quad \forall n \geq \tilde{n}. \quad (8)$$

On one hand, from (8) we immediately deduce that

$$d(x_n, h_n(x_n)) < \eta \quad \forall n \geq \tilde{n}. \quad (9)$$

On the other hand, since $x_n \in \mathcal{GR}(g_n) \setminus U$ then $h_n(x_n) \in \mathcal{GR}(f)$ by inclusion (7). As a consequence

$$d(x_n, h_n(x_n)) \geq \min_{x \in \mathcal{GR}(f)} d(x_n, x) = d(x_n, \mathcal{GR}(f)) \geq \min_{y \in X \setminus U} d(y, \mathcal{GR}(f)) = \eta. \quad (10)$$

Inequalities (9) and (10) provide the required contradiction. \square

Clearly, from the previous proposition, we immediately deduce that if $f \in \text{Hom}(X)$ with $\mathcal{GR}(f) \neq X$ is topologically stable then f does not permit \mathcal{GR} -full explosions. As proved by P. Walters in [16][Theorem 1], any Anosov diffeomorphism on a smooth, compact manifold M without boundary is topologically stable. Consequently, the previous proposition applies in particular to every Anosov diffeomorphism with $\mathcal{GR}(f) \neq M$.

As with Corollary 3.1, the converse of Proposition 3.2 is false in general, as shown in the next example.

Example 3.3. On the circle $\mathbb{S}^1 := \mathbb{R}/2\pi\mathbb{Z}$ embedded in \mathbb{R}^2 with the usual induced topology, for $n \geq 1$ consider the points

$$P_n := \left(\cos\left(\frac{\pi}{2^n}\right), \sin\left(\frac{\pi}{2^n}\right) \right)$$

and

$$Q_n := \left(\cos\left(\frac{\pi}{2^n}\right), -\sin\left(\frac{\pi}{2^n}\right) \right).$$

Let $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ be a homeomorphism which fixes exactly $(1, 0)$ and every P_n and Q_n and such that:

- (i) Every Q_{n-1} and P_n with $n \geq 2$ even is an attractor.
- (ii) Every Q_n and P_{n-1} with $n \geq 2$ even is a repeller.




Figure 4: The dynamical system of Example 3.3.

We refer to Figure 4. In such a case,

$$Fix(f) = \mathcal{GR}(f) = \mathcal{CR}(f).$$

On one hand, since $Fix(f)$ is an infinite set, f is not topologically stable (see [20][Theorem 1]). On the other hand, since $\mathcal{GR}(f) = \mathcal{CR}(f)$, f does not permit \mathcal{GR} -explosions (see Corollary 3.1). (It is also easy to verify directly that f does not permit \mathcal{GR} -explosions.)

We conclude this section by discussing a sufficient condition to avoid \mathcal{GR} -full explosions. This condition is an immediate corollary of Theorem 2.2.

Proposition 3.3. *Let $f \in Hom(X)$ be such that $\mathcal{GR}(f) \neq X$. If $\mathcal{CR}(f) \neq X$ then f does not permit \mathcal{GR} -full explosions.*

Proof. Arguing by contradiction, we assume that f admits \mathcal{GR} -full explosions. This means that there exists a sequence $(g_n)_{n \in \mathbb{N}}$ in $Hom(X)$ converging to f in the uniform topology such that $\mathcal{GR}(g_n) = X$ for any $n \in \mathbb{N}$. Consequently, $\mathcal{GR}(g_n) = \mathcal{CR}(g_n) = X$ for any $n \in \mathbb{N}$. Since by hypothesis $\mathcal{CR}(f) \neq X$, we conclude that f permits \mathcal{CR} -explosions and this fact contradicts Theorem 2.2. \square

Note that, in general, the converse of Proposition 3.3 may be false: in Examples 3.1 and 3.2 we have defined homeomorphisms on a compact metric space such that $\mathcal{GR}(f) \subsetneq \mathcal{CR}(f) = X$ and f does not permit \mathcal{GR} -explosions. In particular, f does not permit \mathcal{GR} -full explosions.

In the following final example, we slightly modify Example 3.1 in order to obtain $f \in Hom(X)$ such that $\mathcal{GR}(f) \subsetneq \mathcal{CR}(f) = X$ and f does not permit \mathcal{GR} -full explosions even though f permits \mathcal{GR} -explosions.




Figure 5: The dynamical system of Example 3.4.

Example 3.4. Referring to the compact metric space X constructed in the previous Example 3.1, let $a, b \in \mathbb{S}^1 \setminus K$ be such that the closed interval $[a, b] \subset \mathbb{S}^1$ (obtained by connecting a and b clockwise) does not intersect K . Let

$$X' = X \cup X_{a,b}$$

where $X_{a,b}$ is an arc (disjoint from X) from a to b .

Define the homeomorphism $f : X' \rightarrow X'$ as follows. f fixes a, b, K , every copy of the interval I and the arc $X_{a,b}$, while f moves all the points in $\mathbb{S}^1 \setminus (K \cup \{a, b\})$ counterclockwise (see Figure 5).

In this case

$$\mathcal{CR}(f) = X' \quad \text{and} \quad \mathcal{GR}(f) = \text{Fix}(f).$$

Essentially the same argument as in Example 3.1 shows that f does not permit \mathcal{GR} -full explosions. However, f does permit \mathcal{GR} -explosions. Indeed, for any $\varepsilon > 0$ there exists $g \in \text{Hom}(X')$ such that

$$d_{\mathcal{C}^0}(f, g) < \varepsilon$$

and g modifies the dynamics so that on the arc $X_{a,b}$, points move from a to b and thus points of the open interval (a, b) can return to themselves with arbitrary strong chains. This means that

$$\mathcal{GR}(g) = \mathcal{GR}(f) \cup (a, b).$$

4 \mathcal{GR} -explosions and cycles

The goal of this section is to explain the relations between explosions and “cycles” for the generalized recurrent set. Let $f \in \text{Hom}(X)$ be such that $\mathcal{GR}(f) \neq X$. We start by introducing the notions of decomposition and cycle for $\mathcal{GR}(f)$.

Given a compact, invariant set $L \subseteq X$, we define by

$$W^s(L) := \{x \in X : d(f^k(x), L) \rightarrow 0 \text{ as } k \rightarrow +\infty\}$$

and

$$W^u(L) := \{x \in X : d(f^{-k}(x), L) \rightarrow 0 \text{ as } k \rightarrow +\infty\}$$

the stable and unstable set, respectively, of L for f . In particular, we have that $x \in W^s(L)$ if and only if $\omega(x) \subseteq L$ and $x \in W^u(L)$ if and only if $\alpha(x) \subseteq L$.

A decomposition of $\mathcal{GR}(f)$ is a finite family L_1, \dots, L_k of compact, invariant, pairwise disjoint sets in X such that

$$\mathcal{GR}(f) = \bigcup_{i=1}^k L_i.$$

Definition 4.1. Let L_1, \dots, L_k be a decomposition of $\mathcal{GR}(f)$.

(i) Let $i, j \in \{1, \dots, k\}$. We write $L_i \rightarrow L_j$ if

$$(W^u(L_i) \cap W^s(L_j)) \setminus \mathcal{GR}(f) \neq \emptyset.$$

Equivalently, there is a point $x \in X$ outside $\mathcal{GR}(f)$ whose orbit is going from L_i to L_j .

(ii) We say that L_{i_1}, \dots, L_{i_r} form an $r \geq 1$ cycle of $\{L_i\}$ if

$$L_{i_1} \rightarrow L_{i_2} \rightarrow \dots \rightarrow L_{i_r} \rightarrow L_{i_1}.$$

(iii) The decomposition $\{L_i\}$ has no cycles if no subset of $\{L_i\}$ forms an $r \geq 1$ cycle.

We first establish that the existence of a decomposition of $\mathcal{GR}(f)$ with no cycles prohibits \mathcal{GR} -explosions. This theorem generalizes the corresponding result –due to Pugh and Shub– for the non-wandering set (see Theorem 6.1 in [14] for flows and Theorem 5.6 in [18] for homeomorphisms). Since in the proof we apply the \mathcal{C}^0 closing lemma, the ambient space is a compact, topological manifold M with $\dim(M) \geq 2$.

Theorem 4.1. Let $f \in \text{Hom}(M)$ be defined on a compact topological manifold M of $\dim(M) \geq 2$ and such that $\mathcal{GR}(f) \neq M$. If there exists a decomposition of $\mathcal{GR}(f)$ without cycles, then f does not permit \mathcal{GR} -explosions.

The proof of Theorem 4.1 uses the same techniques of Pugh-Shub and is postponed to Appendix A. In order to establish the converse of Theorem 4.1, we need to introduce the notion of cycle for a so-called open decomposition of $f \in \text{Hom}(X)$ with $\mathcal{GR}(f) \neq X$.

Given an open set $V \subset X$, we define

$$V^u := \{x \in X : \exists m \geq 0 \text{ such that } f^{-m}(x) \in V\} = \bigcup_{m \geq 0} f^m(V). \quad (11)$$

An open decomposition of f is a finite family V_1, \dots, V_k of open sets in X , with pairwise disjoint closures, such that

$$\mathcal{GR}(f) \subseteq \bigcup_{i=1}^k V_i.$$

Definition 4.2. Let V_1, \dots, V_k be an open decomposition of f .

(i) Let $i, j \in \{1, \dots, k\}$, $i \neq j$. We write $V_i \geq V_j$ if

$$V_j \cap V_i^u \neq \emptyset.$$

Equivalently, there are $x \in V_j$ and $m \geq 0$ ¹ such that $f^{-m}(x) \in V_i$.

(ii) We say that V_{i_1}, \dots, V_{i_r} form an $r > 1$ cycle of $\{V_i\}$ if

$$V_{i_1} \geq V_{i_2} \geq \dots \geq V_{i_r} \geq V_{i_1}.$$

We say that V_j forms a 1 cycle of $\{V_i\}$ if there are $x \notin V_j$ and $m, q > 0$ such that

$$f^m(x), f^{-q}(x) \in V_j.$$

(iii) The open decomposition $\{V_i\}$ has no cycles if no subset of $\{V_i\}$ forms an $r \geq 1$ cycle.

By using the above formalism, we prove the other implication of Theorem 4.1.

Theorem 4.2. Let $f \in \text{Hom}(M)$ be defined on a compact topological manifold M of $\dim(M) \geq 2$ and such that $\mathcal{GR}(f) \neq M$. If f does not permit \mathcal{GR} -explosions, then there exists a decomposition of $\mathcal{GR}(f)$ with no cycles.

The above theorem is a straightforward consequence of the next one.

Theorem 4.3. Let $f \in \text{Hom}(M)$ be defined on a compact topological manifold M of $\dim(M) \geq 2$ and such that $\mathcal{GR}(f) \neq M$. If f does not permit \mathcal{GR} -explosions, then for any open neighborhood U of $\mathcal{GR}(f)$ there exists an open decomposition V_1, \dots, V_k of f with no cycles such that

$$\mathcal{GR}(f) \subset \bigcup_{i=1}^k V_i \subseteq U.$$

Since the proof of Theorem 4.3 follows the same lines of the proof of Lemma 2 in [15], it is postponed to Appendix B.

Proof of Theorem 4.2. Arguing by contradiction, we suppose that every decomposition L_1, \dots, L_k admits an $r \geq 1$ cycle

$$L_{i_1} \rightarrow L_{i_2} \rightarrow \dots \rightarrow L_{i_r} \rightarrow L_{i_1}.$$

This means that there are $x_1, \dots, x_r \notin \mathcal{GR}(f)$ such that

$$\alpha(x_j) \subseteq L_{i_j}, \quad \omega(x_j) \subseteq L_{i_{j+1}} \quad \forall j \in \{1, \dots, r-1\}$$

and

$$\alpha(x_r) \subseteq L_{i_r}, \quad \omega(x_r) \subseteq L_{i_1}.$$

¹Actually, $m > 0$ since the open sets $\{V_i\}$ have pairwise disjoint closures.

Since the $\{L_i\}$ are compact and pairwise disjoint, we can choose a family $\{\mathcal{L}_i\}$ of open sets, with pairwise disjoint closures such that $L_i \subset \mathcal{L}_i$ for any $i = 1, \dots, k$. Let us consider the open neighborhood

$$U := \bigcup_{i=1}^k \mathcal{L}_i.$$

of $\mathcal{GR}(f)$. By shrinking each \mathcal{L}_i a bit, we can assume that the points $x_1, \dots, x_r \notin U$. Then, by construction, any open decomposition $\{V_i\}$ of f such that $\mathcal{GR}(f) \subset \bigcup_i V_i \subseteq U$ has cycles. This fact contradicts Theorem 4.3 applied to U and concludes the proof. \square

Finally, Theorems 4.1 and 4.2 give us the equivalence between no \mathcal{GR} -explosions and the existence of a decomposition of $\mathcal{GR}(f)$ with no cycles.

Theorem 4.4. *Let $f \in \text{Hom}(M)$ be defined on a compact topological manifold M of $\dim(M) \geq 2$ and such that $\mathcal{GR}(f) \neq M$. f does not permit \mathcal{GR} -explosions if and only if there exists a decomposition of $\mathcal{GR}(f)$ with no cycles.*

Remark 4.1. We notice that in the proof of the previous theorem –see Appendixes A and B– we do not use the full definition of the generalized recurrent set, but only the fact that $\mathcal{GR}(f)$ is a compact, invariant set, which contains $\mathcal{NW}(f)$ and is contained in $\mathcal{CR}(f)$. Consequently, with the same proof, we can obtain the corresponding results both for $\mathcal{SCR}(f)$ and $\mathcal{CR}(f)$.

Theorem 4.5. *Let $f \in \text{Hom}(M)$ be defined on a compact topological manifold M of $\dim(M) \geq 2$ and such that $\mathcal{SCR}(f) \neq M$. f does not permit \mathcal{SCR} -explosions if and only if there exists a decomposition of $\mathcal{SCR}(f)$ with no cycles.*

Remark 4.2. The corresponding result holds for $\mathcal{CR}(f)$; since \mathcal{CR} -explosions cannot occur (see Theorem 2.2), we see that for $f \in \text{Hom}(M)$ defined on a compact topological manifold M of $\dim(M) \geq 2$, there exists a decomposition of $\mathcal{CR}(f)$ with no cycles. In fact, this holds for compact metric spaces, as a consequence of the existence of a complete Lyapunov function.

5 Applications to strict Lyapunov functions and genericity

Let X be a compact metric space and $f \in \text{Hom}(X)$. We say that a continuous Lyapunov function $u : X \rightarrow \mathbb{R}$ for f is strict if it is not a first integral. Equivalently, this means that $\mathcal{N}(u) \neq X$, i.e. there exists $x \in X$ such that

$$u(f(x)) < u(x).$$

By Theorem 2.1, f admits a continuous strict Lyapunov function if and only if $\mathcal{GR}(f) \neq X$. In this section, we collect the results of Sections 3 and 4 in order to give conditions to give an affirmative answer to this question:

For a given homeomorphism, is the property of admitting a continuous strict Lyapunov function stable under \mathcal{C}^0 perturbations?

From Corollary 3.1, Proposition 3.2 and Proposition 3.3, we deduce the following

Proposition 5.1. *Let $f \in \text{Hom}(X)$ admit a continuous strict Lyapunov function. Suppose that one of these hypotheses holds:*

- (i) $\mathcal{GR}(f) = \mathcal{CR}(f)$;
- (ii) $\mathcal{CR}(f)$ is strictly contained in X ;
- (iii) f is topologically stable.

Then there exists a neighborhood V of f in $\text{Hom}(X)$ such that any $g \in V$ admits a continuous strict Lyapunov function.

Moreover, from Theorem 4.4, we obtain that a sufficient condition is the existence of a decomposition of $\mathcal{GR}(f)$ without cycles.

Proposition 5.2. *Let $f \in \text{Hom}(M)$ admit a continuous strict Lyapunov function. If there exists a decomposition of $\mathcal{GR}(f)$ without cycles then there exists a neighborhood V of f in $\text{Hom}(M)$ such that any $g \in V$ admits a continuous strict Lyapunov function.*

We finally remark that the property of admitting a continuous strict Lyapunov function is generic. For this purpose, let M be a smooth, compact manifold with metric d . We endow $\text{Hom}(M)$ with the metric

$$d_{\mathcal{H}}(f, g) = \max_{x \in M} \left(\max [d(f(x), g(x)), d(f^{-1}(x), g^{-1}(x))] \right).$$

With this metric, $\text{Hom}(M)$ is a complete space and therefore it is a Baire space. A property in $\text{Hom}(M)$ is said to be generic if the set of $f \in \text{Hom}(M)$ satisfying this property contains a residual set, i.e. a countable intersection of open dense sets.

Proposition 5.3. *On a smooth, compact manifold M , the property in $\text{Hom}(M)$ of admitting a continuous strict Lyapunov function is generic.*

Proof. By Theorem 3.1 in [9], the property in $\text{Hom}(M)$ of having $\text{int}(\mathcal{CR}(f)) = \emptyset$ is generic and so also that of having $\mathcal{GR}(f) \neq M$. Therefore, by Theorem 2.1, the property in $\text{Hom}(M)$ of admitting a continuous strict Lyapunov function is generic too. \square

A Proof of Theorem 4.1

Arguing by contradiction, we suppose that there are an open neighborhood U of $\mathcal{GR}(f)$ in M , a sequence $(h_n)_{n \in \mathbb{N}} \in \text{Hom}(M)$ converging to f in the uniform topology and a sequence of points $(a_n)_{n \in \mathbb{N}} \in \mathcal{GR}(h_n) \setminus U$. Since X is compact and U is open, we can assume that $a_n \rightarrow b_1 \notin \mathcal{GR}(f)$. By assumption

$$a_n \in \mathcal{GR}(h_n) \subseteq \mathcal{CR}(h_n)$$

and therefore –by the \mathcal{C}^0 closing lemma– for each $n \in \mathbb{N}$ there exists $g_n \in \text{Hom}(M)$ such that

$$d_{\mathcal{C}^0}(h_n, g_n) < \frac{1}{n} \quad \text{and} \quad a_n \in \text{Per}(g_n).$$

Denote by $T_n \geq 1$ the least period of a_n and define

$$k_n := nT_n.$$

Clearly, $k_n \rightarrow +\infty$ and

$$g_n^{k_n}(a_n) = a_n \quad \forall n \in \mathbb{N}.$$

This means that there exist $g_n \rightarrow f$ in the \mathcal{C}^0 topology and $k_n \rightarrow +\infty$ such that every point a_n can be equivalently represented by $g_n^{k_n}(a_n)$. Recall that

$$g_n^{k_n}(a_n) = a_n \rightarrow b_1 \notin \mathcal{GR}(f).$$

The alpha limit and the omega limit of every point of M are contained in $\mathcal{NW}(f) \subseteq \mathcal{GR}(f)$ (see Proposition 3.3.4 in [10]). In particular, $\alpha(b_1), \omega(b_1) \subseteq L_1 \cup \dots \cup L_k$ and therefore, by Theorem 5.4 in [18], there exist $i_0, i_1 \in \{1, \dots, k\}$ such that $\alpha(b_1) \subseteq L_{i_0}$ and $\omega(b_1) \subseteq L_{i_1}$ (equivalently $b_1 \in W^u(L_{i_0}) \cap W^s(L_{i_1})$). Since we have no cycles $i_0 \neq i_1$.

Take a compact neighborhood U_{i_1} of L_{i_1} such that

$$b_1 \notin U_{i_1} \quad \text{and} \quad U_{i_1} \cap L_j = \emptyset \quad \forall j \neq i_1.$$

Equivalently, since the L_j are invariant,

$$b_1 \notin U_{i_1} \quad \text{and} \quad f(U_{i_1}) \cap L_j = \emptyset \quad \forall j \neq i_1.$$

As in [18][Page 147], we denote pieces of g_n -orbits from a_n to itself as:

$$\begin{aligned} [a_n, a_n] &:= (a_n, g_n(a_n), g_n^2(a_n), \dots, g_n^{k_n}(a_n)), \\ [a_n, a_n] &:= (a_n, g_n(a_n), g_n^2(a_n), \dots, g_n^{k_n-1}(a_n)), \\ (a_n, a_n) &:= (g_n(a_n), g_n^2(a_n), \dots, g_n^{k_n-1}(a_n)). \end{aligned}$$

Since $a_n \rightarrow b_1 \in W^s(L_{i_1})$, $g_n \rightarrow f$ in the uniform topology and $k_n \rightarrow +\infty$, there exists a sequence of points $p_n \in [a_n, a_n]$ such that

$$d(p_n, L_{i_1}) \rightarrow 0, \quad \text{for } n \rightarrow +\infty.$$

In particular, since $b_1 \notin U_{i_1}$, $p_n \in (a_n, a_n)$ for large n . Moreover, for the same reason, for n sufficiently large, there is $m_n \geq 1$ such that

$$p_n, g_n(p_n), \dots, g_n^{m_n-1}(p_n) \in \text{int } U_{i_1}$$

and

$$g_n^{m_n}(p_n) \notin \text{int } U_{i_1}.$$

Then

$$z_n := g_n^{m_n}(p_n) \in g_n(\text{int } U_{i_1}) \setminus \text{int } U_{i_1} \subset g_n(U_{i_1}) \setminus \text{int } U_{i_1}.$$

Note that, since $d(p_n, L_{i_1}) \rightarrow 0$, $g_n \rightarrow f$ in the \mathcal{C}^0 topology and L_{i_1} is f -invariant, we have

$$m_n \rightarrow +\infty.$$

Finally, since $g_n \rightarrow f$ in the uniform norm, we have that for n large enough

$$g_n(U_{i_1}) \cap L_j = \emptyset \quad \forall j \neq i_1.$$

Let b_2 be a limit point of the sequence $(z_n)_{n \in \mathbb{N}}$. Hence $b_2 \notin \mathcal{GR}(f)$. Moreover, since the first m_n iterates of z_n with respect to g_n^{-1} are contained in U_{i_1} , $g_n \rightarrow f$ in the uniform norm and $m_n \rightarrow +\infty$, it follows that $f^{-m}(b_2) \in U_{i_1}$ for all $m \geq 1$ (recall that, since X is compact, it holds also that $g_n^{-1} \rightarrow f^{-1}$ in the \mathcal{C}^0 topology). This means that $\alpha(b_2) \subseteq L_{i_1}$ or equivalently $b_2 \in W^u(L_{i_1})$. Moreover, from the hypothesis that there are no cycles, $\omega(b_2) \subseteq L_{i_2}$ with $i_2 \neq i_0, i_1$.

Recall now that $z_n \in (a_n, a_n]$. This means that there exists $r_n \geq 1$ such that z_n (which is the first point of the g_n -orbit of p_n outside $\text{int } U_{i_1}$) can be represented as

$$z_n = g_n^{r_n}(a_n).$$

In order to proceed similarly with $b_2 \notin \mathcal{GR}(f)$, we need to prove that also the sequence

$$k_n - r_n \rightarrow +\infty.$$

Suppose to the contrary that $k_n - r_n$ is uniformly bounded. This means that

$$b_1 = \lim_{n \rightarrow +\infty} g_n^{k_n - r_n}(z_n)$$

has the same alpha limit of $b_2 = \lim_{n \rightarrow +\infty} z_n$, that is $\alpha(b_1) \subseteq L_{i_1}$. Since $\alpha(b_1) \subseteq L_{i_0}$ with $i_1 \neq i_0$, this is the desired contradiction. In particular, $z_n \in (a_n, a_n)$ for large n .

We now apply the same argument to $b_2 \notin \mathcal{GR}(f)$. Take a compact neighborhood U_{i_2} of L_{i_2} such that

$$b_2 \notin U_{i_2} \quad \text{and} \quad f(U_{i_2}) \cap L_j = \emptyset \quad \forall j \neq i_2.$$

In order to continue, we denote pieces of g_n -orbits from $z_n = g_n^{r_n}(a_n)$ to a_n as:

$$[z_n, a_n] := (z_n, g_n(z_n), g_n^2(z_n), \dots, g_n^{k_n - r_n}(z_n)),$$

$$[z_n, a_n] := (z_n, g_n(z_n), g_n^2(z_n), \dots, g_n^{k_n - r_n - 1}(z_n)),$$

$$(z_n, a_n) := (g_n(z_n), g_n^2(z_n), \dots, g_n^{k_n - r_n - 1}(z_n)).$$

As in the previous case, there exists a sequence of points $p'_n \in [z_n, a_n]$ such that

$$d(p'_n, L_{i_2}) \rightarrow 0, \quad \text{for } n \rightarrow +\infty.$$

In particular, since $b_2 \notin U_{i_2}$, $p'_n \in (z_n, a_n)$ for large n . Moreover, for n sufficiently large, there is $m'_n \geq 1$ such that

$$p'_n, g_n(p'_n), \dots, g_n^{m'_n-1}(p'_n) \in \text{int } U_{i_2}$$

and

$$g_n^{m'_n}(p'_n) \notin \text{int } U_{i_2}.$$

Then

$$z'_n := g_n^{m'_n}(p'_n) \in g_n(\text{int } U_{i_2}) \setminus \text{int } U_{i_2} \subset g_n(U_{i_2}) \setminus \text{int } U_{i_2}$$

and

$$m'_n \rightarrow +\infty.$$

Finally, since $g_n \rightarrow f$ in the uniform norm, we have that for n large enough

$$g_n(U_{i_2}) \cap L_j = \emptyset \quad \forall j \neq i_2.$$

Let b_3 be a limit point of the sequence $(z'_n)_{n \in \mathbb{N}}$. Hence, $b_3 \notin \mathcal{GR}(f)$ and $\alpha(b_3) \subseteq L_{i_2}$. Moreover, since there are no cycles, $\omega(b_3) \subseteq L_{i_3}$ with $i_3 \neq i_0, i_1, i_2$.

We finally notice that there exists $r'_n \geq 1$ such that z'_n can be represented as

$$z'_n = g_n^{r'_n}(a_n).$$

Arguing as for $k_n - r_n$, it can be shown that the sequence $k_n - r'_n \rightarrow +\infty$. In particular, $z'_n \in (z_n, a_n)$ for large n .

We can proceed with b_3 exactly as b_2 and produce, by iteration, a chain of L_j 's with no repetitions and length greater than k . This is the desired contradiction and the theorem is proved. \square

B Proof of Theorem 4.3

We begin with a definition and a technical lemma.

Definition B.1. Let $f \in \text{Hom}(X)$, $y, z \in X$ and $\varepsilon > 0$. A \mathcal{GR} -chain of ε balls of f from y to z is a finite sequence $\{B_i\} = B_1, \dots, B_n$ of open convex balls such that

- (i) $y \in B_1$ and $z \in B_n$;
- (ii) For any $i \in \{1, \dots, n\}$, $\text{diam}(B_i) < \varepsilon$ and $B_i \cap \mathcal{GR}(f) \neq \emptyset$;
- (iii) For any $i \in \{1, \dots, n-1\}$, there exists $m(i) \geq 0$ such that $f^{m(i)}(B_i) \cap B_{i+1} \neq \emptyset$.

The same definition of chain of balls is given in [15][Page 590] for the non-wandering set. Moreover, Lemma B.1 below is a generalization of Lemma 3 in [15].

Lemma B.1. Let $f \in \text{Hom}(M)$ be defined on a compact topological manifold M of $\dim(M) \geq 2$. Let $y, z \in M$ and let $\varepsilon > 0$. Given a \mathcal{GR} -chain of ε balls of f from y to z , there exists $g \in \text{Hom}(M)$ such that:

- (a) $d_{\mathcal{C}^0}(f, g) < 4\pi\varepsilon$;
- (b) $g^N(f^{-1}(y)) = f(z)$ for some $N > 0$;
- (c) $g = f$ outside the union of the ε balls of the \mathcal{GR} -chain.

Proof of Lemma B.1. Let $\{B_i\} = B_1, \dots, B_n$ be a \mathcal{GR} -chain of ε balls from y to z . For any $i \in \{1, \dots, n-1\}$, let $m(i) \geq 0$ be the first integer such that $f^{m(i)}(B_i) \cap B_{i+1} \neq \emptyset$. For any $i \in \{1, \dots, n-1\}$, choose

$$w_{i+1} \in f^{m(i)}(B_i) \cap B_{i+1}$$

and denote

$$z_i := f^{-m(i)}(w_{i+1}) \in B_i$$

Moreover, since $\{B_i\}$ is a \mathcal{GR} -chain, for any $i \in \{1, \dots, n\}$ we can select

$$k_i \in B_i \cap \mathcal{GR}(f).$$

In particular, each $k_i \in \mathcal{CR}(f)$. Consequently, let $(r_{i,1} = k_i, r_{i,2}, \dots, r_{i,N(k_i)} = k_i)$ be a ε -chain from $r_{i,1} = k_i$ to itself.

Let now consider the following couples of points:

$$(y, k_1)$$

$$(f(r_{1,j}), r_{1,j+1}) \quad \forall j = 1, \dots, N(k_1) - 2$$

$$(f(r_{1,N(k_1)-1}), z_1)$$

$$(f^j(z_1), f^j(z_1)) \quad \forall j = 1, \dots, m(1) - 1$$

$$(f^{m(1)}(z_1), k_2) = (w_2, k_2)$$

$$(f(r_{2,j}), r_{2,j+1}) \quad \forall j = 1, \dots, N(k_2) - 2$$

$$(f(r_{2,N(k_2)-1}), z_2)$$

$$(f^j(z_2), f^j(z_2)) \quad \forall j = 1, \dots, m(2) - 1$$

and so on, until

$$(f^{m(n-1)}(z_{n-1}), z) = (w_n, z)$$

$$(f(z), f(z)).$$

By perturbing f a little, we can assume that for every (q_i, p_i) , (q_j, p_j) with $i \neq j$, it holds that $q_i \neq q_j$ and $p_i \neq p_j$. Moreover, by construction, for any couple (q, p) given above, the distance $d(q, p) < 2\epsilon$. Consequently, applying Lemma 13 in [12], we obtain a homeomorphism $\eta : M \rightarrow M$ such that

$$d_{\mathcal{C}^0}(\eta \circ f, f) < 4\pi\epsilon \quad \text{and} \quad \eta(q) = p$$

for any such a couple (q, p) .

We finally prove that there exists $N > 0$ such that $(\eta \circ f)^N(f^{-1}(y)) = f(z)$. Indeed

$$(\eta \circ f)(f^{-1}(y)) = k_1, \quad (\eta \circ f)(k_1) = r_{1,2}, \quad (\eta \circ f)(r_{1,2}) = r_{1,3}, \quad \dots \quad (\eta \circ f)(r_{1,N(k_1)-1}) = z_1$$

$$(\eta \circ f)(z_1) = f(z_1), \quad (\eta \circ f)(f(z_1)) = f^2(z_1), \quad \dots \quad (\eta \circ f)(f^{m(1)-1}(z_1)) = k_2$$

and so on, until

$$(\eta \circ f)(f^{m(n-1)-1}(z_{n-1})) = \eta(w_n) = z, \quad \eta \circ f(z) = f(z).$$

This proves that, after a number $N > 0$ of iterations of $\eta \circ f$, we have

$$(\eta \circ f)^N(f^{-1}(y)) = f(z)$$

and so $g := \eta \circ f$ is the desired perturbation. \square

We finally prove Theorem 4.3 by using the same techniques of Lemma 2 in [15].

Proof of Theorem 4.3. Let U be an arbitrary open neighborhood of $\mathcal{GR}(f)$ in M . Since f does not permit \mathcal{GR} -explosions, there exists $\epsilon > 0$ such that if the \mathcal{C}^0 distance from $g \in \text{Hom}(M)$ to f is less than $4\pi\epsilon$ then

$$\mathcal{GR}(g) \subset U. \quad (12)$$

Let $\{B_\alpha\}$ be a finite covering of $\mathcal{GR}(f)$ by open convex balls such that, for any α ,

$$B_\alpha \subseteq U \quad \text{and} \quad \text{diam}(B_\alpha) < \epsilon.$$

We denote by $\{U_\beta\}$ the connected components of $\bigcup_\alpha B_\alpha$. By shrinking the balls $\{B_\alpha\}$ if necessary, we can assume that the $\{U_\beta\}$ have pairwise disjoint closures. We now introduce the following equivalence relation on pairs of $\{U_\beta\}$. We say that U_{β_1} is related to U_{β_2} if either $U_{\beta_1} = U_{\beta_2}$ or there is a common cycle containing both U_{β_1} and U_{β_2} , according to Definition 4.2. Moreover, we indicate by \tilde{U}_i the union

of the members of the same relation class. By construction, $\{\tilde{U}_i\}$ is an open decomposition of f with no r cycles for $r > 1$. In order to obtain an open decomposition of f with no r cycles for $r \geq 1$, we define

$$\tilde{U}_i^s := \{x \in X : \exists q \geq 0 \text{ such that } f^q(x) \in \tilde{U}_i\}$$

and (see also formula (11))

$$V_i := \tilde{U}_i^s \cap \tilde{U}_i^u.$$

Then V_1, \dots, V_k is an open decomposition of f with no cycles. That is, $\{V_i\}$ is a finite family of open sets with pairwise disjoint closures such that $\mathcal{GR}(f) \subseteq \bigcup_{i=1}^k V_i$ and no subset of $\{V_i\}$ forms an $r \geq 1$ cycle.

It remains to prove that $V_i \subseteq U$ for any i . Let i be fixed. Arguing by contradiction, we suppose there exists

$$x \in V_i \setminus U.$$

Since $V_i = \tilde{U}_i^s \cap \tilde{U}_i^u$, there exist $m, q > 0$ such that

$$y := f^m(x) \in \tilde{U}_i \quad \text{and} \quad z := f^{-q}(x) \in \tilde{U}_i$$

Let $m, q > 0$ be the minimal integers with this property. Recall that, by definition, \tilde{U}_i is the union of the members of $\{B_\alpha\}$ in the same relation class. This means that there exists a \mathcal{GR} -chain of ε balls of f from y to z and therefore we can apply Lemma B.1. Let $g \in \text{Hom}(M)$ be the homeomorphism given by Lemma B.1. Since $m, q > 0$ are the minimal integers such that $y = f^m(x) \in \tilde{U}_i$ and $z = f^{-q}(x) \in \tilde{U}_i$, the points

$$f^i(x) \quad \forall i \in \{0, \dots, m-1\}$$

and

$$f^{-j}(x) \quad \forall j \in \{0, \dots, q-1\}$$

are outside the union of the ε balls of the \mathcal{GR} -chain. Consequently, by point (iii) of Lemma B.1,

$$g(f^i(x)) = f(f^i(x)) \quad \forall i \in \{0, \dots, m-1\} \tag{13}$$

and

$$g(f^{-j}(x)) = f(f^{-j}(x)) \quad \forall j \in \{0, \dots, q-1\}.$$

Since $x = f^{-m}(y) = f^q(z)$, the previous equalities become respectively

$$g(f^{-i}(y)) = f(f^{-i}(y)) \quad \forall i \in \{1, \dots, m\}$$

and

$$g(f^j(z)) = f(f^j(z)) \quad \forall j \in \{1, \dots, q\}. \tag{14}$$

Consequently, by (14):

$$x = f^q(z) = f(f^{q-1}(z)) = g(f^{q-1}(z)) = g^2(f^{q-2}(z)) = \dots = g^{q-1}(f(z)).$$

Moreover, by point (ii) of Lemma B.1, $g^N(f^{-1}(y)) = f(z)$ for some $N > 0$ and therefore (see also (13))

$$\begin{aligned} x &= g^{N+q-1}(f^{-1}(y)) = g^{N+q-1}(f^{m-1}(x)) = g^{N+q-1}(f(f^{m-2}(x))) = g^{N+q}(f^{m-2}(x)) \\ &= \dots = g^{N+q+m-2}(x). \end{aligned}$$

This means that $x \in \text{Per}(g)$.

Recall that, by property (i) of Lemma B.1, $g \in \text{Hom}(M)$ is such that $d_{\mathcal{G}^0}(f, g) < 4\pi\varepsilon$. Consequently, by (12), $\mathcal{GR}(g) \subset U$. Since $\text{Per}(g) \subseteq \mathcal{GR}(g)$, the point x should belong to U and this gives us the required contradiction. \square

References

- [1] Akin, Ethan. The General Topology of Dynamical Systems (Graduate Studies in Mathematics, 1). American Mathematical Society, Providence, RI, (1993).
- [2] Akin, Ethan; Auslander, Joseph. Generalized recurrence, compactifications, and the Lyapunov topology. *Studia Math.* 201, no. 1, 49–63, (2010).

- [3] Auslander, Joseph. Generalized recurrence in dynamical systems. *Contributions to Differential Equations*, vol. 3, 65–74, (1964).
- [4] Block, Louis; Franke, John E. The chain recurrent set, attractors, and explosions. *Ergodic Theory Dynam. Systems* 5, no. 3, 321–327, (1985).
- [5] De Vries, Jan. Topological dynamical systems. An introduction to the dynamics of continuous mappings. *De Gruyter Studies in Mathematics*, 59. De Gruyter, Berlin, (2014).
- [6] Easton, Robert. Chain transitivity and the domain of influence of an invariant set. *The structure of attractors in dynamical systems*, pp. 95–102, *Lecture Notes in Math.*, 668, Springer, Berlin, (1978).
- [7] Fathi, Albert; Pageault, Pierre. Aubry-Mather theory for homeomorphisms. *Ergodic Theory Dynam. Systems* 35, no. 4, 1187–1207, (2015).
- [8] Franks, John. A variation on the Poincaré-Birkhoff theorem. *Hamiltonian dynamical systems* (Boulder, CO, 1987), 111–117, *Contemp. Math.*, 81, Amer. Math. Soc., Providence, RI, (1988).
- [9] Hurley, Mike. Properties of attractors of generic homeomorphisms. *Ergodic Theory Dynam. Systems* 16, no. 6, 1297–1310, (1996).
- [10] Katok, Anatole; Hasselblatt, Boris. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, (1995).
- [11] Nitecki, Zbigniew. *Differentiable dynamics. An introduction to the orbit structure of diffeomorphisms*. The M.I.T. Press, Cambridge, Mass.-London, xv+282 pp. (1971).
- [12] Nitecki, Zbigniew; Shub, Michael. Filtrations, decompositions, and explosions. *Amer. J. Math.* 97 (1975), no. 4, 1029–1047.
- [13] Palis, Jacob. Ω -explosions. *Proc. Amer. Math. Soc.* 27, 85–90, (1971).
- [14] Pugh, Charles; Shub, Michael. The Ω -Stability Theorem for Flows. *Invent. Math.* 11, 150–158, (1970).
- [15] Shub, Michael; Smale, Steve. Beyond hyperbolicity. *Annals of Mathematics* 96, no. 3, 576–591, (1972).
- [16] Walters, Peter. Anosov diffeomorphisms are topologically stable. *Topology* 9, 71–78, (1970).
- [17] Walters, Peter. On the pseudo-orbit tracing property and its relationship to stability. *The structure of attractors in dynamical systems* (Proc. Conf., North Dakota State Univ., Fargo, N.D., 1977), pp. 231–244, *Lecture Notes in Math.*, 668, Springer, Berlin, (1978).
- [18] Wen, Lan. *Differentiable dynamical systems. An introduction to structural stability and hyperbolicity*. Graduate Studies in Mathematics, 173. American Mathematical Society, Providence, RI, xi+192 pp. (2016).
- [19] Wiseman, Jim. The generalized recurrent set and strong chain recurrence. *Ergodic Theory Dynam. Systems* 38, no. 2, 788–800, (2018).
- [20] Yano, Koichi. Topologically stable homeomorphisms on the circle. *Nagoya Math. J.* 79, 145–149, (1980).
- [21] Yokoi, Katsuya. On strong chain recurrence for maps. *Annales Polonici Mathematici* 114, no. 2, 165–177, (2015).