



ITEs2007 Sixth Italian-Spanish conference
on General Topology and applications

Bressanone, 26-29 June 2007

Sequential order of compact sequential spaces. II

Chiara Baldovino

Università degli Studi di Milano

Dipartimento di Matematica “Federigo Enriques”, Via C. Saldini 50, 20133 Milano, Italy

`baldo@mat.unimi.it`

Baškirov proved, under CH, that there are compact Hausdorff spaces of any sequential order up to and including ω_1 . The results of Baškirov, concisely presented in a Doklady article, are completely revisited and compared with the results of Dow under MA. It is proved that if K_α and K_β are Baškirov spaces of sequential order α and β respectively, then $K_\alpha \times K_\beta$ has sequential order $\max(\alpha, \beta)$.

References

- [1] A. I. Baškirov, *The classification of quotient maps and sequential bicomacta*, Soviet Math. Dokl. **15** (1974), 1104–1109.
- [2] A. Dow, *On mad families and sequential order*, see <http://www.math.uncc.edu/~adow/Others.html> under the title “On the sequential order of Compact spaces”.
- [3] ———, *Sequential order under ma*, Topology Appl. **146-147** (2005), 501–510.
- [4] V. Kannan, *Ordinal invariants in topology II. Sequential order of compactifications*, Compos. Math. **39** (1979), no. 2, 247–262.
- [5] T. Nogura and A. Shibakov, *Sequential order of product spaces*, Topology Appl. **65** (1995), 271–285.